

thm_2Earithmetic_2EMOD_EQ_0
(TMEgNQ36zCEzPUo3byeMgfbJnkTPgBHYrGH)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 1 We define c_2Emin_2E3D to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o(x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define c_2Ebool_2E21 to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (ap (ap (c_2Emin_2E3D (2^{A-27a})) (\lambda V1x \in 2.V1x)) (\lambda V2x \in 2.V2x)))$

Definition 4 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum. (ap c_2Enum_2EABS_num (c_2Enum_2EREP_num m))$

Let $c_2Earithmetic_2EDIV : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EDIV \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{5}$$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{6}$$

Definition 5 We define c_2Enum_2E0 to be $(ap c_2Enum_2EABS_num c_2Enum_2EZERO_REP)$.

Let $c_2Earithmetic_2EMOD : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EMOD \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (7)$$

Definition 6 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21) 2) (\lambda V0t \in 2.V0t)$.

Definition 7 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow P \Rightarrow Q)$ of type ι .

Definition 8 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF))$

Definition 9 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21) 2) (\lambda V2t \in 2.V2t)))$

Definition 10 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge P x))$ of type $\iota \Rightarrow \iota$.

Definition 11 We define $c_2Ebool_2E_3F$ to be $\lambda A.^{2^A}.\lambda P \in (2^A)^{2^A}.(ap V0P (ap (c_2Emin_2E_40)))$

Definition 12 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.$

Let $c_2Earithmetic_2E_2A : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2A \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (8)$$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum. \\ & ((ap (ap c_2Earithmetic_2E_2B c_2Enum_2E0) V0m) = V0m) \wedge (((ap (\\ & ap c_2Earithmetic_2E_2B V0m) c_2Enum_2E0) = V0m) \wedge (((ap (ap c_2Earithmetic_2E_2B \\ & (ap c_2Enum_2ESUC V0m)) V1n) = (ap c_2Enum_2ESUC (ap (ap c_2Earithmetic_2E_2B \\ & V0m) V1n))) \wedge ((ap (ap c_2Earithmetic_2E_2B V0m) (ap c_2Enum_2ESUC \\ & V1n)) = (ap c_2Enum_2ESUC (ap (ap c_2Earithmetic_2E_2B V0m) V1n)))))))))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & (\forall V0n \in ty_2Enum_2Enum.((p (ap (ap c_2Eprim_rec_2E_3C \\ & c_2Enum_2E0) V0n)) \Rightarrow (\forall V1k \in ty_2Enum_2Enum.((V1k = (ap (\\ & ap c_2Earithmetic_2E_2B (ap (ap c_2Earithmetic_2E_2A (ap (ap c_2Earithmetic_2EDIV \\ & V1k) V0n)) V0n)) (ap (ap c_2Earithmetic_2EMOD V1k) V0n))) \wedge (p (ap \\ & (ap c_2Eprim_rec_2E_3C (ap (ap c_2Earithmetic_2EMOD V1k) V0n))))))))) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned}
& (\forall V0n \in ty_2Enum_2Enum. (\forall V1k \in ty_2Enum_2Enum. (\\
& \quad \forall V2r \in ty_2Enum_2Enum. ((\exists V3q \in ty_2Enum_2Enum. (\\
& (V1k = (ap (ap c_2Earithmetic_2E_2B (ap (ap c_2Earithmetic_2E_2A \\
& \quad V3q) V0n)) V2r)) \wedge (p (ap (ap c_2Eprim_rec_2E_3C V2r) V0n)))) \Rightarrow (\\
& \quad (ap (ap c_2Earithmetic_2EMOD V1k) V0n) = V2r))))))
\end{aligned} \tag{12}$$

Assume the following.

$$True \tag{13}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \tag{14}$$

Theorem 1

$$\begin{aligned}
& (\forall V0n \in ty_2Enum_2Enum. ((p (ap (ap c_2Eprim_rec_2E_3C \\
& c_2Enum_2E0) V0n)) \Rightarrow (\forall V1k \in ty_2Enum_2Enum. ((ap (ap c_2Earithmetic_2EMOD \\
& (ap (ap c_2Earithmetic_2E_2A V1k) V0n)) V0n) = c_2Enum_2E0))))
\end{aligned}$$