

thm_2Earithmetic_2EMOD__P__UNIV
(TMJniGcB6EB96PHCX7Lg6uSmoQi2U5WJbji)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Earithmetic_2EDIV : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EDIV \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Earithmetic_2EMOD : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EMOD \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{5}$$

Definition 3 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda \tau a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1Q \in 2.V1Q))\ (\lambda V2R \in 2.V2R))\ (\lambda V3S \in 2.V3S))$

Definition 5 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 6 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow Q)$ of type ι .

Definition 7 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E_3D_3D_3E\ V0t)\ c_2Ebool_2EF))$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \wedge True) \Leftrightarrow \\
& (p \ V0t)) \wedge (((False \wedge (p \ V0t)) \Leftrightarrow False) \wedge (((p \ V0t) \wedge False) \Leftrightarrow False) \wedge \\
& (((p \ V0t) \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)))))) \quad (14)
\end{aligned}$$

Assume the following.

$$\forall A.27a.nonempty \ A.27a \Rightarrow (\forall V0x \in A.27a.((V0x = V0x) \Leftrightarrow True)) \quad (15)$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \Leftrightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow True) \Leftrightarrow \\
& (p \ V0t)) \wedge (((False \Leftrightarrow (p \ V0t)) \Leftrightarrow \neg(p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow False) \Leftrightarrow \neg(\\
& p \ V0t)))))) \quad (16)
\end{aligned}$$

Theorem 1

$$\begin{aligned}
& (\forall V0P \in (2^{ty_2Enum_2Enum}).(\forall V1m \in ty_2Enum_2Enum. \\
& (\forall V2n \in ty_2Enum_2Enum.((p \ (ap \ (ap \ c.2Eprim_rec.2E_3C \\
c.2Enum_2E0) \ V2n)) \Rightarrow ((p \ (ap \ V0P \ (ap \ (ap \ c.2Earithmetic.2EMOD \ V1m) \\
V2n))) \Leftrightarrow (\forall V3q \in ty_2Enum_2Enum.(\forall V4r \in ty_2Enum_2Enum. \\
(((V1m = (ap \ (ap \ c.2Earithmetic.2E_2B \ (ap \ (ap \ c.2Earithmetic.2E_2A \\
V3q) \ V2n)) \ V4r)) \wedge (p \ (ap \ (ap \ c.2Eprim_rec.2E_3C \ V4r) \ V2n))) \Rightarrow (p \\
(ap \ V0P \ V4r)))))))))
\end{aligned}$$