

thm_2Earithmetic_2EMULT__COMM (TMS- Bes5TqECfKDDePxBdzpmFw1SSKyA2zon)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Earithmetic_2E_2A : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2A \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{2}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_T$ to be $(ap (ap (c_2Emin_2E_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})))$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & (ap (ap c_2Earithmetic_2E_2A V0m) V1n) = (ap (ap c_2Earithmetic_2E_2A \\ & V1n) V0m)))) \end{aligned} \tag{3}$$

Theorem 1

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & (ap (ap c_2Earithmetic_2E_2A V0m) V1n) = (ap (ap c_2Earithmetic_2E_2A \\ & V1n) V0m)))) \end{aligned}$$