

thm\_2Earithmetic\_2EMULT\_\_EQ\_\_0  
(TMd8vJNz4EggHzwGokinb4etEGWuLm3B3zW)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.$ if  $(\exists x \in A.p (ap P x))$  then (the  $(\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ ).

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c\_2Emin\_2E\_40 A$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{2}$$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{3}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{4}$$

**Definition 4** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

**Definition 5** We define  $c\_2Earithmetic\_2EZERO$  to be  $c\_2Enum\_2E0$ .

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{5}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{6}$$

**Definition 6** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 7** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num (ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 9** We define  $c\_2Earithmetic\_2EBIT1$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap (ap c\_2Earithmetic\_2E\_2B V0m) V1n))$

**Definition 10** We define  $c\_2Earithmetic\_2ENUMERAL$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

Let  $c\_2Earithmetic\_2E\_2A : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2A \in ((ty\_2Enum\_2Enum)^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum} \quad (7)$$

**Definition 11** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 12** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

**Definition 13** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 14** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

**Definition 15** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_5C\_2F V0t))$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum. \\ & ((ap (ap c\_2Earithmetic\_2E\_2B c\_2Enum\_2E0) V0m) = V0m) \wedge (((ap ( \\ & ap c\_2Earithmetic\_2E\_2B V0m) c\_2Enum\_2E0) = V0m) \wedge (((ap (ap c\_2Earithmetic\_2E\_2B \\ & (ap c\_2Enum\_2ESUC V0m)) V1n) = (ap c\_2Enum\_2ESUC (ap (ap c\_2Earithmetic\_2E\_2B \\ & V0m) V1n))) \wedge ((ap (ap c\_2Earithmetic\_2E\_2B V0m) (ap c\_2Enum\_2ESUC \\ & V1n)) = (ap c\_2Enum\_2ESUC (ap (ap c\_2Earithmetic\_2E\_2B V0m) V1n))))))))) \end{aligned} \quad (8)$$

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum.((V0m = c\_2Enum\_2E0) \vee (\exists V1n \in ty\_2Enum\_2Enum.(V0m = (ap c\_2Enum\_2ESUC V1n)))))) \quad (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum. \\ & ((ap (ap c\_2Earithmetic\_2E\_2A c\_2Enum\_2E0) V0m) = c\_2Enum\_2E0) \wedge \\ & (((ap (ap c\_2Earithmetic\_2E\_2A V0m) c\_2Enum\_2E0) = c\_2Enum\_2E0) \wedge \\ & (((ap (ap c\_2Earithmetic\_2E\_2A (ap c\_2Earithmetic\_2ENUMERAL \\ & (ap c\_2Earithmetic\_2EBIT1 c\_2Earithmetic\_2EZERO))) V0m) = V0m) \wedge \\ & (((ap (ap c\_2Earithmetic\_2E\_2A V0m) (ap c\_2Earithmetic\_2ENUMERAL \\ & (ap c\_2Earithmetic\_2EBIT1 c\_2Earithmetic\_2EZERO))) = V0m) \wedge \\ & ((ap (ap c\_2Earithmetic\_2E\_2A (ap c\_2Enum\_2ESUC V0m)) V1n) = (ap \\ & (ap c\_2Earithmetic\_2E\_2B (ap (ap c\_2Earithmetic\_2E\_2A V0m) V1n)) \\ & V1n)) \wedge ((ap (ap c\_2Earithmetic\_2E\_2A V0m) (ap c\_2Enum\_2ESUC V1n)) = \\ & (ap (ap c\_2Earithmetic\_2E\_2B V0m) (ap (ap c\_2Earithmetic\_2E\_2A \\ & V0m) V1n))))))))) \end{aligned} \quad (10)$$

Assume the following.

$$True \quad (11)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (12)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2.(((True \vee (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \vee True) \Leftrightarrow True) \wedge (((False \vee (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee False) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee (p V0t)) \Leftrightarrow (p V0t)))))) \quad (14)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t)) \wedge ((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (15)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0x \in A.27a.((V0x = V0x) \Leftrightarrow True)) \quad (16)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (17)$$

Assume the following.

$$(\forall V0n \in ty\_2Enum\_2Enum.(\neg((ap c\_2Enum\_2ESUC V0n) = c\_2Enum\_2E0))) \quad (18)$$

### Theorem 1

$$(\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.((ap (ap c\_2Earithmetic\_2E\_2A V0m) V1n) = c\_2Enum\_2E0) \Leftrightarrow ((V0m = c\_2Enum\_2E0) \vee (V1n = c\_2Enum\_2E0))))$$