

thm_2Earithmetic_2ENORM_0
(TML2CyAR1z5Nwa3TRRYgPYVYLNuiZbjuweQ)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)$.

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a})))$

Definition 4 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 5 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 6 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Assume the following.

$$True \tag{4}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \tag{5}$$

Theorem 1 $((ap\ c_2Earithmetic_2ENUMERAL\ c_2Earithmetic_2EZERO) = c_2Enum_2E0)$.