

thm_2Earithmetic_2ENOT_NUM_EQ
(TMc3439a7ancC7bNdpV1k1rc6QdE7xy4FvE)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda P \in 2.(ap (ap (c_2Emin_2E_3D (2^{A-27a})) (\lambda V0t \in 2.V0t)) (\lambda V1t \in 2.V1t))$

Definition 4 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num (ap (ap (c_2Emin_2E_3D (2^{A-27a})) (\lambda V0t \in 2.V0t)) (\lambda V1t \in 2.V1t)) m))$

Definition 5 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 6 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 7 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF))$

Definition 8 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)) t1))$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then (the $(\lambda x.x \in A \wedge p x)$) of type $\iota \Rightarrow \iota$.

Definition 10 We define $c_Ebool_2E_3F$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A-27a}). (ap\ V0P\ (ap\ (c_Emin_2E_40$

Definition 11 We define $c_Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Definition 12 We define $c_Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c_Ebool_2E_21\ 2)\ (\lambda V2t \in$

Definition 13 We define $c_Earithmetic_2E_3C_3D$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Assume the following.

$$(\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (V0m = V1n) \Leftrightarrow ((p\ (ap\ (ap\ c_Earithmetic_2E_3C_3D\ V0m)\ V1n)) \wedge (p\ (ap\ (ap\ c_Earithmetic_2E_3C_3D\ V1n)\ V0m)))))) \quad (5)$$

Assume the following.

$$(\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\neg(p\ (ap\ (ap\ c_Earithmetic_2E_3C_3D\ V0m)\ V1n))) \Leftrightarrow (p\ (ap\ (ap\ c_Earithmetic_2E_3C_3D\ (ap\ c_Enum_2ESUC\ V1n))\ V0m)))))) \quad (6)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((p\ V0A) \vee (p\ V1B)) \Leftrightarrow ((p\ V1B) \vee (p\ V0A)))))) \quad (7)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((\neg((p\ V0A) \wedge (p\ V1B))) \Leftrightarrow ((\neg(p\ V0A) \vee \neg(p\ V1B)))) \wedge (((\neg(p\ V0A) \vee (p\ V1B)) \Leftrightarrow ((\neg(p\ V0A) \wedge \neg(p\ V1B)))))))))) \quad (8)$$

Theorem 1

$$(\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\neg(V0m = V1n)) \Leftrightarrow ((p\ (ap\ (ap\ c_Earithmetic_2E_3C_3D\ (ap\ c_Enum_2ESUC\ V0m))\ V1n)) \vee (p\ (ap\ (ap\ c_Earithmetic_2E_3C_3D\ (ap\ c_Enum_2ESUC\ V1n))\ V0m))))))$$