

# thm\_2Earithmetic\_2ENOT\_\_SUC\_\_ADD\_\_LESS\_\_EQ (TMNU6PLPfQAK56XwoEe51Q7uFdDHhdRSnTD)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{2}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{3}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{4}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num ($

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{5}$$

**Definition 5** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 6** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 7** We define `c_2Ebool_2E_7E` to be  $(\lambda V0t \in 2. (ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_7E))$

**Definition 8** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2. (ap (c_2Ebool_2E_21 2) V2t) V1t2)) V0t1)))$

**Definition 9** We define `c_2Emin_2E_40` to be  $\lambda A. \lambda P \in 2^A. \mathbf{if} (\exists x \in A. p (ap P x)) \mathbf{then} (the (\lambda x. x \in A \wedge p (ap P x))) \mathbf{else} (ap P x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define `c_2Ebool_2E_3F` to be  $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap V0P (ap (c_2Emin_2E_40) (ap P x))))$

**Definition 11** We define `c_2Eprim_rec_2E_3C` to be  $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum. (ap (ap (ap c_2Eprim_rec_2E_3C V0m) V1n))$

**Definition 12** We define `c_2Ebool_2E_5C_2F` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2. (ap (c_2Ebool_2E_21 2) V2t) V1t2)) V0t1)))$

**Definition 13** We define `c_2Earithmetic_2E_3C_3D` to be  $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum. (ap (ap (ap c_2Earithmetic_2E_3C_3D V0m) V1n))$

Assume the following.

$$(\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (p (ap (ap (ap c_2Eprim_rec_2E_3C V0m) V1n)) \Leftrightarrow (p (ap (ap c_2Earithmetic_2E_3C_3D (ap c_2Enum_2ESUC V0m)) V1n)))))) \quad (6)$$

Assume the following.

$$(\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (p (ap (ap c_2Earithmetic_2E_3C_3D V0m) (ap (ap c_2Earithmetic_2E_2B V0m) V1n)))))) \quad (7)$$

Assume the following.

$$(\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\neg (p (ap (ap c_2Eprim_rec_2E_3C V0m) V1n))) \Leftrightarrow (p (ap (ap c_2Earithmetic_2E_3C_3D V1n) V0m)))))) \quad (8)$$

Assume the following.

$$True \quad (9)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg (p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg (p V0t))))) \quad (10)$$

**Theorem 1**

$$\neg (p (ap (ap c_2Earithmetic_2E_3C_3D (ap c_2Enum_2ESUC (ap (ap c_2Earithmetic_2E_2B V0m) V1n)) V0m))))$$