

thm_2Earithmetic_2ENUMERAL_MULT_EQ_DIV
 (TMSvsVN-
 vbRvJ6CBr2SsNcUB88waso1MG1Em)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (1)$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap (ap (c_2Emin_2E_3D (2^{A_27a})) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)))$

Definition 4 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum. V0x$.

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (2)$$

Let $c_2Enum_2EAABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EAABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (3)$$

Definition 5 We define c_2Enum_2E0 to be $(ap c_2Enum_2EAABS_num c_2Enum_2EZERO_REP)$.

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (4)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (5)$$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum. (ap c_2Enum_2EAABS_num (c_2Enum_2E0 m))$

Let c_2 be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (6)$$

Definition 7 We define $c_2Earithmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap\ (ap\ c_2Earithmetic_2EBIT1\ n)\ V)$

Definition 8 We define $c_2Earithmetic_2EBIT2$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap (ap c_2Earithmetic_2EBIT2$

Let c_2 be given. Assume the following.

$$c_2Earithmetic_2EMOD \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \\ (7)$$

Let $c_2Earithmetic_2EDIV : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EDIV \in ((ty_2Enum_2Enum ty_2Enum_2Enum) ty_2Enum_2Enum) \\ (8)$$

Let $c_2Earithmetic_2E_2A : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2A \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (9)$$

Definition 9 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 10 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t\ t \in 2.V0t))$.

Definition 11 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_7E))$

Definition 12 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in$

Definition 13 We define c_2 to be $\lambda A. \lambda P \in 2^A$. if $(\exists x \in A. p(ap P x))$ then $(the (\lambda x. x \in A \wedge$

Definition 14 We define $\in \mathbb{E}\text{bool}$ to be $\lambda A. \exists a : \cup. (\lambda V0P \in (2^{A-27a}).(ap\;V0P\;(ap\;(\in\;\mathbb{E}\text{min})\;2E\;40)$

Definition 15. We define \mathcal{C} 2Eprim, rec 2E 3C to be $\lambda V0m \in tu\; 2Enum\; 2Enum\; \lambda V1n \in tu\; 2Enum\; 2Enum$

Assume the following

(VÄ)

Assume the following.

$$\begin{aligned}
 & (\forall V0x \in ty_2Enum_2Enum. (\forall V1y \in ty_2Enum_2Enum. (\forall V2z \in ty_2Enum_2Enum. ((p (ap (ap c_2Eprim_rec_2E_3C c_2Enum_2E0) \\
 & V0x)) \Rightarrow (((ap (ap c_2Earithmetic_2E_2A V0x) V1y) = V2z) \Leftrightarrow ((V1y = \\
 & ap (ap c_2Earithmetic_2EDIV V2z) V0x)) \wedge ((ap (ap c_2Earithmetic_2EMOD \\
 & V2z) V0x) = c_2Enum_2E0))))))) \\
 \end{aligned} \tag{11}$$

Assume the following.

$$True \tag{12}$$

Assume the following.

$$\begin{aligned}
 & (\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\
 & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))))
 \end{aligned} \tag{13}$$

Assume the following.

$$\begin{aligned}
 & (\forall V0n \in ty_2Enum_2Enum. (p (ap (ap c_2Eprim_rec_2E_3C c_2Enum_2E0) \\
 & (ap c_2Enum_2ESUC V0n)))) \\
 \end{aligned} \tag{14}$$

Theorem 1

$$\begin{aligned}
 & (\forall V0x \in ty_2Enum_2Enum. (\forall V1y \in ty_2Enum_2Enum. (\forall V2z \in ty_2Enum_2Enum. (((ap (ap c_2Earithmetic_2E_2A \\
 & (ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT1 V0x))) \\
 & V1y) = (ap c_2Earithmetic_2ENUMERAL V2z)) \Leftrightarrow ((V1y = (ap (ap c_2Earithmetic_2EDIV \\
 & (ap c_2Earithmetic_2ENUMERAL V2z)) (ap c_2Earithmetic_2ENUMERAL \\
 & (ap c_2Earithmetic_2EBIT1 V0x)))) \wedge ((ap (ap c_2Earithmetic_2EMOD \\
 & (ap c_2Earithmetic_2ENUMERAL V2z)) (ap c_2Earithmetic_2ENUMERAL \\
 & (ap c_2Earithmetic_2EBIT1 V0x))) = c_2Enum_2E0))) \wedge (((ap (ap c_2Earithmetic_2E_2A \\
 & (ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT2 V0x))) \\
 & V1y) = (ap c_2Earithmetic_2ENUMERAL V2z)) \Leftrightarrow ((V1y = (ap (ap c_2Earithmetic_2EDIV \\
 & (ap c_2Earithmetic_2ENUMERAL V2z)) (ap c_2Earithmetic_2ENUMERAL \\
 & (ap c_2Earithmetic_2EBIT2 V0x)))) \wedge ((ap (ap c_2Earithmetic_2EMOD \\
 & (ap c_2Earithmetic_2ENUMERAL V2z)) (ap c_2Earithmetic_2ENUMERAL \\
 & (ap c_2Earithmetic_2EBIT2 V0x))) = c_2Enum_2E0)))))))
 \end{aligned}$$