

thm\_2Earithmetic\_2EODD\_EXISTS  
(TMdeS8t2ZQCGyCoiUz8aKSqNF3m9STX1Pdm)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{2}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{3}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{4}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num ($

Let  $c\_2Earithmetic\_2EODD : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2EODD \in (2^{ty\_2Enum\_2Enum}) \tag{5}$$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{6}$$

**Definition 5** We define  $c\_2Enum\_2E0$  to be  $(ap c\_2Enum\_2EABS\_num c\_2Enum\_2EZERO\_REP)$ .

**Definition 6** We define  $c\_2Earithmetic\_2EZERO$  to be  $c\_2Enum\_2E0$ .

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})ty\_2Enum\_2Enum) \quad (7)$$

**Definition 7** We define  $c\_2Earithmetic\_2EBIT2$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap (ap c\_2Earithmetic\_2E\_2B$

**Definition 8** We define  $c\_2Earithmetic\_2ENUMERAL$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

Let  $c\_2Earithmetic\_2E\_2A : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2A \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})ty\_2Enum\_2Enum) \quad (8)$$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\mathbf{if} (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p$   
of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A.\lambda P \in 2^A.(ap V0P \in (2^{A-27a}).(ap V0P (ap (c\_2Emin\_2E\_40$

Let  $c\_2Earithmetic\_2EEVEN : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2EEVEN \in (2^{ty\_2Enum\_2Enum}) \quad (9)$$

**Definition 11** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$   
of type  $\iota$ .

**Definition 12** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 13** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_7E$

**Definition 14** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t$

Assume the following.

$$(\forall V0n \in ty\_2Enum\_2Enum.(p (ap c\_2Earithmetic\_2EODD (ap c\_2Enum\_2ESUC (ap (ap c\_2Earithmetic\_2E\_2A (ap c\_2Earithmetic\_2ENUMERAL (ap c\_2Earithmetic\_2EBIT2 c\_2Earithmetic\_2EZERO))) V0n)))))) \quad (10)$$

Assume the following.

$$(\forall V0n \in ty\_2Enum\_2Enum.(((p (ap c\_2Earithmetic\_2EEVEN V0n)) \Rightarrow (\exists V1m \in ty\_2Enum\_2Enum.(V0n = (ap (ap c\_2Earithmetic\_2E\_2A (ap c\_2Earithmetic\_2ENUMERAL (ap c\_2Earithmetic\_2EBIT2 c\_2Earithmetic\_2EZERO))) V1m)))) \wedge ((p (ap c\_2Earithmetic\_2EODD V0n)) \Rightarrow (\exists V2m \in ty\_2Enum\_2Enum.(V0n = (ap c\_2Enum\_2ESUC (ap (ap c\_2Earithmetic\_2E\_2A (ap c\_2Earithmetic\_2ENUMERAL (ap c\_2Earithmetic\_2EBIT2 c\_2Earithmetic\_2EZERO))) V2m)))))))))) \quad (11)$$

Assume the following.

$$True \quad (12)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(p V0t)))))) \quad (14)$$

**Theorem 1**

$$(\forall V0n \in ty\_2Enum\_2Enum.((p (ap c\_2Earithmetic\_2EODD V0n)) \Leftrightarrow (\exists V1m \in ty\_2Enum\_2Enum.(V0n = (ap c\_2Enum\_2ESUC (ap (ap c\_2Earithmetic\_2E\_2A (ap c\_2Earithmetic\_2ENUMERAL (ap c\_2Earithmetic\_2EBIT2 c\_2Earithmetic\_2EZERO)) V1m)))))))$$