

thm_2Earithmetic_2EOR_LESS
(TMTpu48yZGXkgc9wghkDp4kH4VMuhdh9uth)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 4 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num ($

Definition 5 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 6 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 7 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF$

Definition 8 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then (the $(\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$).

Definition 10 We define $c_Ebool_2E_3F$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ V0P\ (ap\ (c_Emin_2E_40$

Definition 11 We define $c_Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum.$

Definition 12 We define $c_Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c_Ebool_2E_21\ 2)\ (\lambda V2t \in$

Definition 13 We define $c_Earithmetic_2E_3C_3D$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum.$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\ & (p\ (ap\ (ap\ c_Eprim_rec_2E_3C\ V0m)\ V1n)) \Leftrightarrow (p\ (ap\ (ap\ c_Earithmetic_2E_3C_3D \\ & (ap\ c_Enum_2ESUC\ V0m))\ V1n)))) \end{aligned} \quad (5)$$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\forall A_27a. nonempty\ A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_27a. (p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (7)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((\\ & (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg (p\ V0t)))))) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\ & (p\ (ap\ (ap\ c_Earithmetic_2E_3C_3D\ (ap\ c_Enum_2ESUC\ V0m))\ V1n)) \Rightarrow \\ & (p\ (ap\ (ap\ c_Eprim_rec_2E_3C\ V0m)\ V1n)))) \end{aligned}$$