

thm_2Earithmetic_2EPRE__SUB (TMHg3TdoKH6e2vtLTvqM3hToctNS8KcxAQP)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_7E$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P))$

Definition 4 We define $c_2Ebool_2E_7E$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow P \Rightarrow Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_7E))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num (c_2Enum_2ESUC_REP m))$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge P x))$ of type $\iota \Rightarrow \iota$.

Definition 10 We define $c_Ebool_2E_3F$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ V0P\ (ap\ (c_Emin_2E_40$

Definition 11 We define $c_Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Definition 12 We define $c_Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c_Ebool_2E_21\ 2)\ (\lambda V2t \in$

Definition 13 We define $c_Earithmetic_2E_3C_3D$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Let $c_Earithmetic_2E_2D : \iota$ be given. Assume the following.

$$c_Earithmetic_2E_2D \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (5)$$

Definition 14 We define c_Ebool_2ECOND to be $\lambda A_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A_27a. (\lambda V2t2 \in A_27a. ($

Let $c_Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_Enum_2EZERO_REP \in \omega \quad (6)$$

Definition 15 We define c_Enum_2E0 to be $(ap\ c_Enum_2EABS_num\ c_Enum_2EZERO_REP)$.

Definition 16 We define $c_Eprim_rec_2EPRE$ to be $\lambda V0m \in ty_2Enum_2Enum. (ap\ (ap\ (ap\ (c_Ebool_2E$

Assume the following.

$$\begin{aligned} & ((\forall V0m \in ty_2Enum_2Enum. ((ap\ (ap\ c_Earithmetic_2E_2D \\ & c_Enum_2E0)\ V0m) = c_Enum_2E0)) \wedge (\forall V1m \in ty_2Enum_2Enum. \\ & (\forall V2n \in ty_2Enum_2Enum. ((ap\ (ap\ c_Earithmetic_2E_2D\ (\\ & ap\ c_Enum_2ESUC\ V1m))\ V2n) = (ap\ (ap\ (ap\ (c_Ebool_2ECOND\ ty_2Enum_2Enum) \\ & (ap\ (ap\ c_Eprim_rec_2E_3C\ V1m)\ V2n))\ c_Enum_2E0)\ (ap\ c_Enum_2ESUC \\ & (ap\ (ap\ c_Earithmetic_2E_2D\ V1m)\ V2n))))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\ & ((ap\ (ap\ c_Earithmetic_2E_2D\ V0m)\ V1n) = c_Enum_2E0) \Leftrightarrow (p\ (ap\ (\\ & ap\ c_Earithmetic_2E_3C_3D\ V0m)\ V1n)))))) \end{aligned} \quad (8)$$

Assume the following.

$$True \quad (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p \\ & V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \end{aligned} \quad (10)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p\ V0t))) \quad (11)$$

Assume the following.

$$(\forall V0t \in 2.((p V0t) \vee (\neg(p V0t)))) \quad (12)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \vee (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \vee True) \Leftrightarrow True) \wedge \\ & (((False \vee (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee False) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee \\ & (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \quad (13)$$

Assume the following.

$$\forall A_27a.nonempty \ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (14)$$

Assume the following.

$$\forall A_27a.nonempty \ A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (15)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(\\ & p V0t)))))) \end{aligned} \quad (16)$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty \ A_27a \Rightarrow (\forall V0t1 \in A_27a.(\forall V1t2 \in \\ & A_27a.(((ap (ap (ap (c_2Ebool_2ECOND A_27a) c_2Ebool_2ET) V0t1) \\ & V1t2) = V0t1) \wedge ((ap (ap (ap (c_2Ebool_2ECOND A_27a) c_2Ebool_2EF \\ & V0t1) V1t2) = V1t2)))))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} & (\forall V0P \in (2^{ty_2Enum_2Enum}).(((p (ap V0P c_2Enum_2E0)) \wedge \\ & (\forall V1n \in ty_2Enum_2Enum.((p (ap V0P V1n)) \Rightarrow (p (ap V0P (ap c_2Enum_2ESUC \\ & V1n)))))) \Rightarrow (\forall V2n \in ty_2Enum_2Enum.(p (ap V0P V2n)))))) \end{aligned} \quad (18)$$

Assume the following.

$$\begin{aligned} & (((ap c_2Eprim_rec_2EPRE c_2Enum_2E0) = c_2Enum_2E0) \wedge (\forall V0m \in \\ & ty_2Enum_2Enum.((ap c_2Eprim_rec_2EPRE (ap c_2Enum_2ESUC V0m)) = \\ & V0m))) \end{aligned} \quad (19)$$

Theorem 1

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & (ap c_2Eprim_rec_2EPRE (ap (ap c_2Earithmetic_2E_2D V0m) V1n)) = \\ & (ap (ap c_2Earithmetic_2E_2D (ap c_2Eprim_rec_2EPRE V0m)) V1n)))) \end{aligned}$$