

thm_2Earithmetic_2ESUB__LEFT__GREATER (TMacc4p3G1pWcLBgvdL1NYMdkFUEmEwcKW9)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num (c_2Enum_2ESUC_REP m))$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge P x))$ of type $\iota \Rightarrow \iota$.

Definition 10 We define $c_Ebool_2E_3F$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ V0P\ (ap\ (c_Emin_2E_40$

Definition 11 We define $c_Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Definition 12 We define $c_Earithmetic_2E_3E$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{5}$$

Definition 13 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Let $c_2Earithmetic_2E_2D : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2D \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{7}$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\ & (ap\ (ap\ c_2Earithmetic_2E_2B\ V0m)\ V1n) = (ap\ (ap\ c_2Earithmetic_2E_2B \\ & \quad V1n)\ V0m)))) \end{aligned} \tag{8}$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\ & \quad \forall V2p \in ty_2Enum_2Enum. ((p\ (ap\ (ap\ c_Eprim_rec_2E_3C\ (\\ & \quad ap\ (ap\ c_2Earithmetic_2E_2D\ V0m)\ V1n))\ V2p)) \Leftrightarrow ((p\ (ap\ (ap\ c_Eprim_rec_2E_3C \\ & \quad V0m)\ (ap\ (ap\ c_2Earithmetic_2E_2B\ V1n)\ V2p))) \wedge (p\ (ap\ (ap\ c_Eprim_rec_2E_3C \\ & \quad \quad c_2Enum_2E0)\ V2p)))))) \end{aligned} \tag{9}$$

Assume the following.

$$True \tag{10}$$

Assume the following.

$$\forall A_27a. nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \tag{11}$$

Theorem 1

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\ & \quad \forall V2p \in ty_2Enum_2Enum. ((p\ (ap\ (ap\ c_2Earithmetic_2E_3E \\ & \quad V0m)\ (ap\ (ap\ c_2Earithmetic_2E_2D\ V1n)\ V2p))) \Leftrightarrow ((p\ (ap\ (ap\ c_2Earithmetic_2E_3E \\ & \quad (ap\ (ap\ c_2Earithmetic_2E_2B\ V0m)\ V2p))\ V1n)) \wedge (p\ (ap\ (ap\ c_2Earithmetic_2E_3E \\ & \quad \quad V0m)\ c_2Enum_2E0)))))) \end{aligned}$$