

thm_2Earithmetic_2ESUB__LEFT__LESS (TMQ- vafbXc91mRfN7jdZGDPy6GG2X4XZ9hhJ)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 4 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num ($

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{5}$$

Let $c_2Earithmetic_2E_2D : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2D \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 5 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 6 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 7 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_7E))$

Definition 8 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2)))$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then (the $(\lambda x.x \in A \wedge p)$ of type $\iota \Rightarrow \iota$).

Definition 10 We define $c_2Ebool_2E_3F$ to be $\lambda A.^{27a} : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40$

Definition 11 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum$

Definition 12 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2)))$

Definition 13 We define $c_2Earithmetic_2E_3C_3D$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{7}$$

Definition 14 We define c_2Enum_2E0 to be $(ap c_2Enum_2EABS_num c_2Enum_2EZERO_REP)$.

Assume the following.

$$\begin{aligned} & ((\forall V0n \in ty_2Enum_2Enum.((ap (ap c_2Earithmetic_2E_2B \\ & c_2Enum_2E0) V0n) = V0n)) \wedge (\forall V1m \in ty_2Enum_2Enum.(\forall V2n \in \\ & ty_2Enum_2Enum.((ap (ap c_2Earithmetic_2E_2B (ap c_2Enum_2ESUC \\ & V1m)) V2n) = (ap c_2Enum_2ESUC (ap (ap c_2Earithmetic_2E_2B V1m) \\ & V2n)))))) \end{aligned} \tag{8}$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & (p (ap (ap c_2Eprim_rec_2E_3C V0m) V1n)) \Leftrightarrow (p (ap (ap c_2Earithmetic_2E_3C_3D \\ & (ap c_2Enum_2ESUC V0m)) V1n)))) \end{aligned} \tag{9}$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & \forall V2p \in ty_2Enum_2Enum.((p (ap (ap c_2Earithmetic_2E_3C_3D \\ & V0m) (ap (ap c_2Earithmetic_2E_2D V1n) V2p))) \Leftrightarrow ((p (ap (ap c_2Earithmetic_2E_3C_3D \\ & (ap (ap c_2Earithmetic_2E_2B V0m) V2p)) V1n)) \vee (p (ap (ap c_2Earithmetic_2E_3C_3D \\ & V0m) c_2Enum_2E0)))))) \end{aligned} \tag{10}$$

Assume the following.

$$True \tag{11}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (12)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2.(((True \vee (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \vee True) \Leftrightarrow True) \wedge (((False \vee (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee False) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee (p V0t)) \Leftrightarrow (p V0t)))))) \quad (14)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0x \in A.27a.((V0x = V0x) \Leftrightarrow True)) \quad (15)$$

Assume the following.

$$(\forall V0n \in ty_2Enum_2Enum.(\neg (p (ap (ap c.2Eprim_rec.2E.3C V0n) c.2Enum_2E0)))) \quad (16)$$

Theorem 1

$$(\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\forall V2p \in ty_2Enum_2Enum.((p (ap (ap c.2Eprim_rec.2E.3C V0m) (ap (ap c.2Earithmetic.2E.2D V1n) V2p))) \Leftrightarrow (p (ap (ap c.2Eprim_rec.2E.3C (ap (ap c.2Earithmetic.2E.2B V0m) V2p)) V1n))))))$$