

# thm\_2Earithmetic\_2ESUB\_\_LESS\_\_EQ (TMGr- woMk92Myikmbiq7jNyHjsPGtT9kWmcL)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{2}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{3}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{4}$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num (c\_2Enum\_2ESUC\_REP m))$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge P x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A-27a}). (ap\ V0P\ (ap\ (c\_Emin\_2E\_40$

**Definition 11** We define  $c\_Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum$

**Definition 12** We define  $c\_Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c\_Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

**Definition 13** We define  $c\_Earithmic\_2E\_3C\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{5}$$

**Definition 14** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{6}$$

Let  $c\_2Earithmetic\_2E\_2D : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2D \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{7}$$

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ V1n) = (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V1n)\ V0m)))) \tag{8}$$

Assume the following.

$$p\ (ap\ (ap\ c\_2Earithmetic\_2E\_3C\_3D\ V0m)\ (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ V1n)))) \tag{9}$$

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. ((ap\ (ap\ c\_2Earithmetic\_2E\_2D\ V0m)\ V1n) = c\_2Enum\_2E0) \Leftrightarrow (p\ (ap\ (ap\ c\_2Earithmetic\_2E\_3C\_3D\ V0m)\ V1n)))))) \tag{10}$$

Assume the following.

$$(\forall V0a \in ty\_2Enum\_2Enum. (\forall V1b \in ty\_2Enum\_2Enum. (\forall V2c \in ty\_2Enum\_2Enum. ((ap\ (ap\ c\_2Earithmetic\_2E\_2D\ V0a)\ (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V1b)\ V2c)) = (ap\ (ap\ c\_2Earithmetic\_2E\_2D\ (ap\ (ap\ c\_2Earithmetic\_2E\_2D\ V0a)\ V1b))\ V2c)))))) \tag{11}$$

Assume the following.

$$True \tag{12}$$

Assume the following.

$$\forall A_{27a}. \text{nonempty } A_{27a} \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_{27a}. (p \ V0t)) \Leftrightarrow (p \ V0t))) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow True) \Leftrightarrow (p \ V0t)) \wedge (((False \Leftrightarrow (p \ V0t)) \Leftrightarrow \neg(p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p \ V0t)))))) \quad (14)$$

**Theorem 1**

$$(\forall V0n \in \text{ty\_2Enum\_2Enum}. (\forall V1m \in \text{ty\_2Enum\_2Enum}. (p \ (ap \ (ap \ c\_2Earithmetic\_2E\_3C\_3D \ (ap \ (ap \ c\_2Earithmetic\_2E\_2D \ V0n) \ V1m)) \ V0n))))$$