

thm\_2Earithmetic\_2ESUB\_\_RIGHT\_\_GREATER  
(TMaCjws1kJN46CT4frFqzW6rbZGFu7GcB4X)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A. \lambda x \in A. \lambda y \in A. inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})) (\lambda V1t \in 2.V1t)) (\lambda V2t \in 2.V2t)))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2. \lambda Q \in 2. inj\_o (p \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2. (ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (1)$$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \quad (2)$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \quad (3)$$

Let  $c\_2Enum\_2EAbs\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EAbs\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (4)$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. (ap c\_2Enum\_2EAbs\_num m)$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (ap P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 12** We define  $c\_2Earithmetic\_2E\_3E$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum.$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (5)$$

Let  $c\_2Earithmetic\_2E\_2D : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2D \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (6)$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. ( \\ & (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ V1n) = (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V1n)\ V0m)))) \\ \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. ( \\ & \forall V2p \in ty\_2Enum\_2Enum. ((p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C\ V0m)\ \\ & (ap\ (ap\ c\_2Earithmetic\_2E\_2D\ V1n)\ V2p))) \Leftrightarrow (p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C\ \\ & (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ V2p))\ V1n))))))) \\ \end{aligned} \quad (8)$$

Assume the following.

$$True \quad (9)$$

Assume the following.

$$\forall A\_27a.\ nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (10)$$

### Theorem 1

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. ( \\ & \forall V2p \in ty\_2Enum\_2Enum. ((p\ (ap\ (ap\ c\_2Earithmetic\_2E\_3E\ \\ & (ap\ (ap\ c\_2Earithmetic\_2E\_2D\ V0m)\ V1n))\ V2p)) \Leftrightarrow (p\ (ap\ (ap\ c\_2Earithmetic\_2E\_3E\ \\ & V0m)\ (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V1n)\ V2p))))))) \\ \end{aligned}$$