

thm_2Earithmetic_2ESUB_RIGHT_LESS (TMM3nQk13dDVhdAfPePNU4EnQqZkBmSdM65)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num (ap c_2Enum_2EREP_num (ap c_2Enum_2ESUC_REP m)))$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 10 We define $c_Ebool_2E_3F$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A-27a}). (ap\ V0P\ (ap\ (c_Emin_2E_40$

Definition 11 We define $c_Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{5}$$

Definition 12 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Let $c_2Earithmetic_2E_2D : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2D \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{7}$$

Definition 13 We define $c_Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c_Ebool_2E_21\ 2)\ (\lambda V2t \in$

Definition 14 We define $c_2Earithmetic_2E_3C_3D$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\ & (ap\ (ap\ c_2Earithmetic_2E_2B\ V0m)\ V1n) = (ap\ (ap\ c_2Earithmetic_2E_2B \\ & \quad V1n)\ V0m)))) \end{aligned} \tag{8}$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\ & (\neg(p\ (ap\ (ap\ c_Eprim_rec_2E_3C\ V0m)\ V1n))) \Leftrightarrow (p\ (ap\ (ap\ c_2Earithmetic_2E_3C_3D \\ & \quad V1n)\ V0m)))) \end{aligned} \tag{9}$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\ & \quad \forall V2p \in ty_2Enum_2Enum. ((p\ (ap\ (ap\ c_2Earithmetic_2E_3C_3D \\ & V0m)\ (ap\ (ap\ c_2Earithmetic_2E_2D\ V1n)\ V2p))) \Leftrightarrow ((p\ (ap\ (ap\ c_2Earithmetic_2E_3C_3D \\ & \quad (ap\ (ap\ c_2Earithmetic_2E_2B\ V0m)\ V2p))\ V1n) \vee (p\ (ap\ (ap\ c_2Earithmetic_2E_3C_3D \\ & \quad V0m)\ c_2Enum_2E0)))))) \end{aligned} \tag{10}$$

Assume the following.

$$True \tag{11}$$

Assume the following.

$$\begin{aligned} & ((\forall V0t \in 2. ((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t)) \wedge (((\neg True) \Leftrightarrow False) \wedge \\ & \quad ((\neg False) \Leftrightarrow True))) \end{aligned} \tag{12}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \quad (14)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((\neg((p\ V0A) \wedge (p\ V1B))) \Leftrightarrow ((\neg(p\ V0A) \vee \neg(p\ V1B))) \wedge ((\neg((p\ V0A) \vee (p\ V1B))) \Leftrightarrow ((\neg(p\ V0A) \wedge \neg(p\ V1B)))))))) \quad (15)$$

Theorem 1

$$(\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\forall V2p \in ty_2Enum_2Enum. ((p\ (ap\ (ap\ c_2Eprim_rec_2E_3C\ (ap\ (ap\ c_2Earithmetic_2E_2D\ V0m)\ V1n))\ V2p)) \Leftrightarrow ((p\ (ap\ (ap\ c_2Eprim_rec_2E_3C\ V0m)\ (ap\ (ap\ c_2Earithmetic_2E_2B\ V1n)\ V2p))) \wedge (p\ (ap\ (ap\ c_2Eprim_rec_2E_3C\ c_2Enum_2E0)\ V2p)))))))$$