

thm\_2Earithmetic\_2ESUC\_\_ONE\_\_ADD  
(TMGr85mgEYBMmm6xkSDdKoQL2WArWWbsQJ6)

October 26, 2020

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{1}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{2}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{3}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

**Definition 3** We define  $c\_2Earithmetic\_2EZERO$  to be  $c\_2Enum\_2E0$ .

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{4}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{5}$$

**Definition 4** We define  $c\_2Ebool\_2ET$  to be  $(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 5** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V1x \in 2.V1x))$

**Definition 6** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num\ (c\_2Enum\_2EREP\_num\ (c\_2Enum\_2ESUC\_REP\ m)))$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{6}$$

**Definition 7** We define `c_2Earithmetic_2EBIT1` to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap (ap c\_2Earithmetic\_2EBIT1$

**Definition 8** We define `c_2Earithmetic_2ENUMERAL` to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.(ap (ap c\_2Earithmetic\_2E\_2B V0m) V1n) = (ap (ap c\_2Earithmetic\_2E\_2B V1n) V0m)))) \quad (7)$$

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum.((ap c\_2Enum\_2ESUC V0m) = (ap (ap c\_2Earithmetic\_2E\_2B V0m) (ap c\_2Earithmetic\_2ENUMERAL (ap c\_2Earithmetic\_2EBIT1 c\_2Earithmetic\_2EZERO)))))) \quad (8)$$

**Theorem 1**

$$(\forall V0n \in ty\_2Enum\_2Enum.((ap c\_2Enum\_2ESUC V0n) = (ap (ap c\_2Earithmetic\_2E\_2B (ap c\_2Earithmetic\_2ENUMERAL (ap c\_2Earithmetic\_2EBIT1 c\_2Earithmetic\_2EZERO))) V0n)))$$