

thm\_2Earithmetic\_2ZERO\_\_LESS\_\_ADD  
(TMVs7FEoqM1VRnuTpZY9zxPH7iamNi4Sq2w)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{2}$$

**Definition 1** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.$ if  $(\exists x \in A.p (ap\ P\ x))$  **then** (the  $(\lambda x.x \in A \wedge p)$  of type  $\iota \Rightarrow \iota$ ).

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o\ (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40\ A\ a))))$

**Definition 4** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o\ (p\ P \Rightarrow p\ Q)$  of type  $\iota$ .

**Definition 5** We define  $c\_2Ebool\_2E\_T$  to be  $(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 6** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^{A-27a}))\ P)\ a)))$

**Definition 7** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in 2.V2t))))$

**Definition 8** We define  $c\_2Ebool\_2E\_F$  to be  $(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V0t \in 2.V0t))$ .

**Definition 9** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in 2.V2t))))$

**Definition 10** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap\ (ap\ c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E\_F))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{3}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{omega}) \quad (4)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{omega}) \quad (5)$$

**Definition 11** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. (ap\ c\_2Enum\_2EABS\_num$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (6)$$

**Definition 12** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

**Definition 13** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. ( \\ & ((ap\ (ap\ c\_2Earithmetic\_2E\_2B\ c\_2Enum\_2E0)\ V0m) = V0m) \wedge ((ap\ ( \\ & ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ c\_2Enum\_2E0) = V0m) \wedge ((ap\ (ap\ c\_2Earithmetic\_2E\_2B \\ & (ap\ c\_2Enum\_2ESUC\ V0m))\ V1n) = (ap\ c\_2Enum\_2ESUC\ (ap\ (ap\ c\_2Earithmetic\_2E\_2B \\ & V0m)\ V1n))) \wedge ((ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ (ap\ c\_2Enum\_2ESUC \\ & V1n)) = (ap\ c\_2Enum\_2ESUC\ (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ V1n)))))) \end{aligned} \quad (7)$$

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum. ((V0m = c\_2Enum\_2E0) \vee (\exists V1n \in ty\_2Enum\_2Enum. (V0m = (ap\ c\_2Enum\_2ESUC\ V1n)))))) \quad (8)$$

Assume the following.

$$True \quad (9)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (10)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p\ V0t))) \quad (11)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge \\ & (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee \\ & (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (12)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \quad (14)$$

Assume the following.

$$(\forall V0n \in ty\_2Enum\_2Enum. (\neg(p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C\ V0n)\ V0n)))) \quad (15)$$

Assume the following.

$$(\forall V0n \in ty\_2Enum\_2Enum. (p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C\ c\_2Enum\_2E0)\ (ap\ c\_2Enum\_2ESUC\ V0n)))) \quad (16)$$

**Theorem 1**

$$(\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. (p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C\ c\_2Enum\_2E0)\ (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ V1n)))) \Leftrightarrow ((p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C\ c\_2Enum\_2E0)\ V0m)) \vee (p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C\ c\_2Enum\_2E0)\ V1n))))))$$