

thm_2Ebag_2EBAG__ALL__DISTINCT__BAG__OF__SET
(TMSZVaoLdfd-
kxv7zyQy11gNJFLTmXJboNXD)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V1x \in 2.V1x))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (c_2Enum_2ESUC_REP\ m))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})ty_2Enum_2Enum) \quad (6)$$

Definition 7 We define $c_2Earithmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmetic_2E_2B))$

Definition 8 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Definition 9 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2)) (\lambda V0t \in 2.V0t)$.

Definition 10 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 11 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_21 2))$

Definition 12 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2)) (\lambda V2t \in 2.V2t)))$

Definition 13 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then $(the (\lambda x.x \in A))$ of type $\iota \Rightarrow \iota$.

Definition 14 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40))))$

Definition 15 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.V1n$

Definition 16 We define $c_2Earithmetic_2E_3E$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.V1n$

Definition 17 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2)) (\lambda V2t \in 2.V2t)))$

Definition 18 We define $c_2Earithmetic_2E_3E_3D$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.V1n$

Definition 19 We define $c_2Ebag_2EBAG_INN$ to be $\lambda A_27a : \iota.\lambda V0e \in A_27a.\lambda V1n \in ty_2Enum_2Enum.V1n$

Definition 20 We define $c_2Ebag_2EBAG_IN$ to be $\lambda A_27a : \iota.\lambda V0e \in A_27a.\lambda V1b \in (ty_2Enum_2Enum^A)$

Definition 21 We define $c_2Ebag_2ESET_OF_BAG$ to be $\lambda A_27a : \iota.\lambda V0b \in (ty_2Enum_2Enum^{A_27a}).(\lambda V1x \in A_27a.V1x)$

Definition 22 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Definition 23 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.V2t2)))$

Definition 24 We define $c_2Ebag_2EBAG_OF_SET$ to be $\lambda A_27a : \iota.\lambda V0P \in (2^{A_27a}).(\lambda V1x \in A_27a.(ap V1x P))$

Definition 25 We define $c_2Earithmetic_2E_3C_3D$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.V1n$

Definition 26 We define $c_2Ebag_2EBAG_ALL_DISTINCT$ to be $\lambda A_27a : \iota.\lambda V0b \in (ty_2Enum_2Enum^A)$

Assume the following.

$$\forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow (\forall V0s \in (2^{A_{.27a}}).((ap\ (c_2Ebag_2ESET_OF_BAG\ A_{.27a})\ (ap\ (c_2Ebag_2EBAG_OF_SET\ A_{.27a})\ V0s)) = V0s)) \quad (7)$$

Assume the following.

$$\forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow (\forall V0b \in (ty_2Enum_2Enum^{A_{.27a}}).((p\ (ap\ (c_2Ebag_2EBAG_ALL_DISTINCT\ A_{.27a})\ V0b)) \Leftrightarrow ((ap\ (c_2Ebag_2EBAG_OF_SET\ A_{.27a})\ (ap\ (c_2Ebag_2ESET_OF_BAG\ A_{.27a})\ V0b)) = V0b))) \quad (8)$$

Assume the following.

$$True \quad (9)$$

Assume the following.

$$\forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow (\forall V0x \in A_{.27a}.((V0x = V0x) \Leftrightarrow True)) \quad (10)$$

Theorem 1

$$\forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow (\forall V0s \in (2^{A_{.27a}}).(p\ (ap\ (c_2Ebag_2EBAG_ALL_DISTINCT\ A_{.27a})\ (ap\ (c_2Ebag_2EBAG_OF_SET\ A_{.27a})\ V0s))))$$