

thm_2Ebag_2EBAG_CHOICE_SING (TMX- oPtC8N3DvxzMnjiBXmXz7Eqfm6wp5w3q)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 7 We define `c_2Earithmic_2EBIT1` to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmic_2EBIT1 V0n) V0n)$.

Definition 8 We define `c_2Earithmic_2ENUMERAL` to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Definition 9 We define `c_2Ebool_2EF` to be $(ap (c_2Ebool_2E21 2) (\lambda V0t \in 2.V0t))$.

Definition 10 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 11 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E21 2) (\lambda V2t \in 2.V2t) V1t2) V0t1))$.

Definition 12 We define `c_2Emin_2E_40` to be $\lambda A.\lambda P \in 2^A.\mathbf{if} (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \Rightarrow P x))$ of type $\iota \Rightarrow \iota$.

Definition 13 We define `c_2Ebool_2ECOND` to be $\lambda A.27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A.27a.(\lambda V2t2 \in A.27a.(ap (c_2Ebool_2E21 2) V2t2) V1t1) V0t))$.

Definition 14 We define `c_2Ebag_2EBAG_INSERT` to be $\lambda A.27a : \iota.\lambda V0e \in A.27a.\lambda V1b \in (ty_2Enum_2Enum A.27a)$.

Let `c_2Ebag_2EBAG_CHOICE` : $\iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow c_2Ebag_2EBAG_CHOICE A.27a \in (A.27a^{(ty_2Enum_2Enum A.27a)}) \quad (7)$$

Definition 15 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E21) V0t)$.

Definition 16 We define `c_2Ebool_2E_3F` to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A.27a}).(ap V0P (ap (c_2Emin_2E_40 V0P) V0P)))$.

Definition 17 We define `c_2Eprim_rec_2E_3C` to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.V0m$.

Definition 18 We define `c_2Earithmic_2E_3E` to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.V0m$.

Definition 19 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E21 2) (\lambda V2t \in 2.V2t) V1t2) V0t1))$.

Definition 20 We define `c_2Earithmic_2E_3E_3D` to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.V0m$.

Definition 21 We define `c_2Ebag_2EBAG_INN` to be $\lambda A.27a : \iota.\lambda V0e \in A.27a.\lambda V1n \in ty_2Enum_2Enum.V0e$.

Definition 22 We define `c_2Ebag_2EBAG_IN` to be $\lambda A.27a : \iota.\lambda V0e \in A.27a.\lambda V1b \in (ty_2Enum_2Enum A.27a)$.

Definition 23 We define `c_2Ecombin_2EK` to be $\lambda A.27a : \iota.\lambda A.27b : \iota.(\lambda V0x \in A.27a.(\lambda V1y \in A.27b.V0x))$.

Definition 24 We define `c_2Ebag_2EEMPTY_BAG` to be $\lambda A.27a : \iota.(ap (c_2Ecombin_2EK ty_2Enum_2Enum A.27a) V0e)$.

Assume the following.

$$\begin{aligned} & \forall A.27a.nonempty A.27a \Rightarrow (\forall V0b \in (ty_2Enum_2Enum A.27a). \\ & (\forall V1e1 \in A.27a.(\forall V2e2 \in A.27a.((p (ap (ap (c_2Ebag_2EBAG_IN A.27a) V1e1) (ap (ap (c_2Ebag_2EBAG_INSERT A.27a) V2e2) V0b))) \Leftrightarrow \\ & ((V1e1 = V2e2) \vee (p (ap (ap (c_2Ebag_2EBAG_IN A.27a) V1e1) V0b))))))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. (\forall V1b \in \\ (ty_2Enum_2Enum^{A_27a}). (\neg((ap\ (ap\ (c_2Ebag_2EBAG_INSERT\ A_27a) \\ V0x)\ V1b) = (c_2Ebag_2EEMPTY_BAG\ A_27a)))))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. (\neg(p\ (ap\ (ap \\ (c_2Ebag_2EBAG_IN\ A_27a)\ V0x)\ (c_2Ebag_2EEMPTY_BAG\ A_27a)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0b \in (ty_2Enum_2Enum^{A_27a}). \\ ((\neg(V0b = (c_2Ebag_2EEMPTY_BAG\ A_27a))) \Rightarrow (p\ (ap\ (ap\ (c_2Ebag_2EBAG_IN \\ A_27a)\ (ap\ (c_2Ebag_2EBAG_CHOICE\ A_27a)\ V0b))\ V0b)))) \end{aligned} \quad (11)$$

Assume the following.

$$True \quad (12)$$

Assume the following.

$$\begin{aligned} (\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p \\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \end{aligned} \quad (13)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p\ V0t))) \quad (14)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge \\ (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee \\ (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((\\ (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \end{aligned} \quad (16)$$

Assume the following.

$$\begin{aligned} ((\forall V0t \in 2. ((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge \\ ((\neg False) \Leftrightarrow True))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow \\ True)) \end{aligned} \quad (18)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. (\forall V1y \in A_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (19)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p\ V0t1) \Rightarrow ((p\ V1t2) \Rightarrow (p\ V2t3))) \Leftrightarrow (((p\ V0t1) \wedge (p\ V1t2)) \Rightarrow (p\ V2t3)))))) \quad (20)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in 2. (\forall V1x_27 \in 2. (\forall V2y \in 2. (\forall V3y_27 \in \\ & 2. (((((p\ V0x) \Leftrightarrow (p\ V1x_27)) \wedge ((p\ V1x_27) \Rightarrow ((p\ V2y) \Leftrightarrow (p\ V3y_27)))) \Rightarrow \\ & (((p\ V0x) \Rightarrow (p\ V2y)) \Leftrightarrow ((p\ V1x_27) \Rightarrow (p\ V3y_27)))))))) \end{aligned} \quad (21)$$

Theorem 1

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((ap\ (c_2Ebag_2EBAG_CHOICE\ A_27a)\ (ap\ (ap\ (c_2Ebag_2EBAG_INSERT\ A_27a)\ V0x)\ (c_2Ebag_2EEMPTY_BAG\ A_27a))) = V0x))$$