

thm_2Ebag_2EBAG_DELETE_BAG_IN_down
(TMMU5VsSvnmPKkufdw2VrXRRjaYNpp6ARH1)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 7 We define `c_2Earithmic_2EBIT1` to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmic_2EBIT1 V0n) V0n)$.

Definition 8 We define `c_2Earithmic_2ENUMERAL` to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Definition 9 We define `c_2Ebool_2EF` to be $(ap (c_2Ebool_2E21 2) (\lambda V0t \in 2.V0t))$.

Definition 10 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 11 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E21) V0t)$.

Definition 12 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E21 2) (\lambda V2t \in 2.V2t) V1t2) V0t1))$.

Definition 13 We define `c_2Emin_2E_40` to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge P x)) \text{ else } (the (\lambda x.x \in A \wedge \neg P x)))$ of type $\iota \Rightarrow \iota$.

Definition 14 We define `c_2Ebool_2E_3F` to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A_27a) V0P)))$.

Definition 15 We define `c_2Eprim_rec_2E_3C` to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.V0m$.

Definition 16 We define `c_2Earithmic_2E_3E` to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.V0m$.

Definition 17 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E21 2) (\lambda V2t \in 2.V2t) V1t2) V0t1))$.

Definition 18 We define `c_2Earithmic_2E_3E_3D` to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.V0m$.

Definition 19 We define `c_2Ebag_2EBAG_INN` to be $\lambda A_27a : \iota.\lambda V0e \in A_27a.\lambda V1n \in ty_2Enum_2Enum.V0e$.

Definition 20 We define `c_2Ebag_2EBAG_IN` to be $\lambda A_27a : \iota.\lambda V0e \in A_27a.\lambda V1b \in (ty_2Enum_2Enum^{A_27a}).V0e$.

Definition 21 We define `c_2Ebool_2ECOND` to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.(ap (c_2Ebool_2E21 2) V2t2) V1t1) V0t))$.

Definition 22 We define `c_2Ebag_2EBAG_INSERT` to be $\lambda A_27a : \iota.\lambda V0e \in A_27a.\lambda V1b \in (ty_2Enum_2Enum^{A_27a}).V0e$.

Definition 23 We define `c_2Ebag_2EBAG_DELETE` to be $\lambda A_27a : \iota.\lambda V0b0 \in (ty_2Enum_2Enum^{A_27a}).\lambda V1b1 \in (ty_2Enum_2Enum^{A_27a}).V0b0$.

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow (\forall V0b \in (ty_2Enum_2Enum^{A_27a}). \\ & (\forall V1e1 \in A_27a.(\forall V2e2 \in A_27a.((p (ap (ap (c_2Ebag_2EBAG_IN A_27a) V1e1) (ap (ap (c_2Ebag_2EBAG_INSERT A_27a) V2e2) V0b))) \Leftrightarrow \\ & ((V1e1 = V2e2) \vee (p (ap (ap (c_2Ebag_2EBAG_IN A_27a) V1e1) V0b))))))) \end{aligned} \quad (7)$$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p V0t))) \quad (10)$$

Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_27a. (p V0t)) \Leftrightarrow (p V0t))) \quad (11)$$

Assume the following.

$$(\forall V0t \in 2. (((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \quad (12)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow False) \Leftrightarrow \neg(p V0t)))))) \quad (13)$$

Assume the following.

$$((\forall V0t \in 2. ((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (14)$$

Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (15)$$

Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow (\forall V0x \in A_27a. (\forall V1y \in A_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (16)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(p V0t)))))) \quad (17)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (\forall V2C \in 2. (((p V0A) \vee (p V1B) \vee (p V2C)) \Leftrightarrow (((p V0A) \vee (p V1B)) \vee (p V2C)))))) \quad (18)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((p V0A) \vee (p V1B)) \Leftrightarrow ((p V1B) \vee (p V0A)))) \quad (19)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(\forall V2C \in 2.(((p V0A) \wedge (p V1B) \vee (p V2C))) \Leftrightarrow (((p V0A) \wedge (p V1B)) \vee ((p V0A) \wedge (p V2C)))))) \quad (20)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((p V0A) \Rightarrow (p V1B)) \Leftrightarrow ((\neg(p V0A)) \vee (p V1B)))) \quad (21)$$

Assume the following.

$$(\forall V0P \in 2.(\forall V1Q \in 2.(\forall V2R \in 2.(((p V0P) \vee (p V1Q)) \Rightarrow (p V2R)) \Leftrightarrow (((p V0P) \Rightarrow (p V2R)) \wedge ((p V1Q) \Rightarrow (p V2R)))))) \quad (22)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (23)$$

Assume the following.

$$2.(((p V0x) \Leftrightarrow (p V1x_{.27})) \wedge ((p V1x_{.27}) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_{.27})))) \Rightarrow 2.(((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_{.27}) \Rightarrow (p V3y_{.27})))) \quad (24)$$

Assume the following.

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0f \in (2^{A_{.27a}}).(\forall V1v \in A_{.27a}.((\forall V2x \in A_{.27a}.((V2x = V1v) \Rightarrow (p (ap V0f V2x)))) \Leftrightarrow (p (ap V0f V1v)))))) \quad (25)$$

Theorem 1

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0b0 \in (ty_2Enum_2Enum^{A_{.27a}}).(\forall V1b \in (ty_2Enum_2Enum^{A_{.27a}}).(\forall V2e1 \in A_{.27a}.(\forall V3e2 \in A_{.27a}.(((p (ap (ap (ap (c_2Ebag_2EBAG_DELETE A_{.27a}) V0b0) V2e1) V1b)) \wedge ((\neg(V2e1 = V3e2)) \wedge (p (ap (ap (c_2Ebag_2EBAG_IN A_{.27a}) V3e2) V0b0)))) \Rightarrow (p (ap (ap (c_2Ebag_2EBAG_IN A_{.27a}) V3e2) V1b))))))$$