

thm_2Ebag_2EBAG_DELETE_EMPTY (TM- RoeMZ5TXApZMZpdCtFh8SaGEFdqqkLM6m)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Ecombin_2EK$ to be $\lambda A.\lambda a : \iota.\lambda A.\lambda b : \iota.(\lambda V0x \in A.\lambda V1y \in A.\lambda V0x)$

Definition 4 We define $c_2Ebag_2EEMPTY_BAG$ to be $\lambda A.\lambda a : \iota.(ap\ (c_2Ecombin_2EK\ ty_2Enum_2Enum))$

Definition 5 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 6 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))))$

Definition 7 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 8 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 9 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E_3D_3D_3E\ V0t)\ c_2Ebool_2EF))$

Definition 10 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (4)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (5)$$

Definition 11 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum. (ap\ c_2Enum_2EABS_num$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (6)$$

Definition 12 We define $c_2Earithmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum. (ap\ (ap\ c_2Earithmetic$

Definition 13 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum. V0x$.

Definition 14 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in$

Definition 15 We define $c_2Emin_2E_40$ to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p\ (ap\ P\ x)) \text{ then } (the\ (\lambda x. x \in A \wedge$
of type $\iota \Rightarrow \iota$.

Definition 16 We define c_2Ebool_2ECOND to be $\lambda A. 27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A. 27a. (\lambda V2t2 \in A. 27a. ($

Definition 17 We define $c_2Ebag_2EBAG_INSERT$ to be $\lambda A. 27a : \iota. \lambda V0e \in A. 27a. \lambda V1b \in (ty_2Enum_2Enum$

Definition 18 We define $c_2Ebag_2EBAG_DELETE$ to be $\lambda A. 27a : \iota. \lambda V0b0 \in (ty_2Enum_2Enum^{A. 27a}). \lambda$

Assume the following.

$$\begin{aligned} & \forall A. 27a. nonempty\ A. 27a \Rightarrow (\forall V0x \in A. 27a. (\forall V1b \in \\ & (ty_2Enum_2Enum^{A. 27a}). (\neg((ap\ (ap\ (c_2Ebag_2EBAG_INSERT\ A. 27a) \\ & V0x)\ V1b) = (c_2Ebag_2EEMPTY_BAG\ A. 27a)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall A. 27a. nonempty\ A. 27a \Rightarrow (\forall V0x \in A. 27a. (\forall V1y \in \\ & A. 27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall A. 27a. nonempty\ A. 27a \Rightarrow (\forall V0e \in A. 27a. (\forall V1b \in \\ & (ty_2Enum_2Enum^{A. 27a}). (\neg(p\ (ap\ (ap\ (ap\ (c_2Ebag_2EBAG_DELETE \\ & A. 27a)\ (c_2Ebag_2EEMPTY_BAG\ A. 27a))\ V0e)\ V1b)))))) \end{aligned}$$