

# thm\_2Ebag\_2EBAG\_\_DIFF\_\_2L (TMQm- CaoFkekH2LKvYUBKkpgrqkBgNZv4CiE)

October 26, 2020

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{1}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$\text{nonempty } ty\_2Enum\_2Enum \tag{2}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{3}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A. \lambda x \in A. \lambda y \in A. inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{4}$$

**Definition 3** We define  $c\_2Ebool\_2ET$  to be  $(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A. \lambda 27a : \iota. (\lambda V0P \in (2^{A-27a}). (ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^{A-27a}))\ (\lambda V1P \in 2.V1P))$

**Definition 5** We define  $c\_2Ebag\_2EBAG\_UNION$  to be  $\lambda A. \lambda 27a : \iota. \lambda V0b \in (ty\_2Enum\_2Enum^{A-27a}). \lambda V1c \in (ty\_2Enum\_2Enum^{A-27a}).$

Let  $c\_2Earithmetic\_2E\_2D : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2D \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{5}$$

**Definition 6** We define  $c\_2Ebag\_2EBAG\_DIFF$  to be  $\lambda A. \lambda 27a : \iota. \lambda V0b1 \in (ty\_2Enum\_2Enum^{A-27a}). \lambda V1b2 \in (ty\_2Enum\_2Enum^{A-27a}).$

**Definition 7** We define  $c\_2Ebool\_2EF$  to be  $(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V0t \in 2.V0t))$ .

**Definition 8** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 9** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_7E))$

**Definition 10** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.))$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.( \\ & \forall V2p \in ty\_2Enum\_2Enum.((ap (ap c\_2Earithmetic\_2E\_2D (ap \\ & (ap c\_2Earithmetic\_2E\_2D V0m) V1n)) V2p) = (ap (ap c\_2Earithmetic\_2E\_2D \\ & V0m) (ap (ap c\_2Earithmetic\_2E\_2B V1n) V2p)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & (\forall V0P \in (2^{ty\_2Enum\_2Enum}).(\forall V1a \in ty\_2Enum\_2Enum. \\ & (\forall V2b \in ty\_2Enum\_2Enum.((p (ap V0P (ap (ap c\_2Earithmetic\_2E\_2D \\ & V1a) V2b))) \Leftrightarrow (\forall V3d \in ty\_2Enum\_2Enum.(((V2b = (ap (ap c\_2Earithmetic\_2E\_2B \\ & V1a) V3d)) \Rightarrow (p (ap V0P c\_2Enum\_2E0))) \wedge ((V1a = (ap (ap c\_2Earithmetic\_2E\_2B \\ & V2b) V3d)) \Rightarrow (p (ap V0P V3d)))))))))) \end{aligned} \quad (7)$$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A\_27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (( \\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \end{aligned} \quad (11)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (12)$$

**Theorem 1**

$$\begin{aligned} & \forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0X \in (ty\_2Enum\_2Enum^{A\_27a}). \\ & (\forall V1Y \in (ty\_2Enum\_2Enum^{A\_27a}).(\forall V2Z \in (ty\_2Enum\_2Enum^{A\_27a}). \\ & ((ap (ap (c\_2Ebag\_2EBAG\_DIFF A\_27a) (ap (ap (c\_2Ebag\_2EBAG\_DIFF \\ & A\_27a) V0X) V1Y)) V2Z) = (ap (ap (c\_2Ebag\_2EBAG\_DIFF A\_27a) V0X) \\ & (ap (ap (c\_2Ebag\_2EBAG\_UNION A\_27a) V1Y) V2Z)))))) \end{aligned}$$