

thm_2Ebag_2EBAG__DIFF__EQ__EMPTY
(TMaQ2YkbsrBq5joqaUFAgqSMv95SDXkX6TG)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Ecombin_2EK$ to be $\lambda A.\lambda a : \iota.\lambda A.\lambda b : \iota.(\lambda V0x \in A.\lambda V1y \in A.\lambda V0x)$

Definition 4 We define $c_2Ebag_2EEMPTY_BAG$ to be $\lambda A.\lambda a : \iota.(ap\ (c_2Ecombin_2EK\ ty_2Enum_2Enum))$

Let $c_2Earithmetic_2E_2D : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2D \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{4}$$

Definition 5 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 6 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a})))$

Definition 7 We define $c_2Ebag_2EBAG_DIFF$ to be $\lambda A.\lambda a : \iota.\lambda V0b1 \in (ty_2Enum_2Enum^{A-27a}).\lambda V1b2 \in (ty_2Enum_2Enum^{A-27a}).\lambda V1b2$

Definition 8 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 9 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (10)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\forall V0f \in (A_27b^{A_27a}). (\forall V1g \in (A_27b^{A_27a}). ((V0f = V1g) \Leftrightarrow (\forall V2x \in A_27a. ((ap\ V0f\ V2x) = (ap\ V1g\ V2x)))))) \quad (11)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\forall V0x \in A_27a. (\forall V1y \in A_27b. ((ap\ (ap\ (c_2Ecombin_2EK\ A_27a\ A_27b)\ V0x)\ V1y) = V0x))) \quad (12)$$

Theorem 1

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0b \in (ty_2Enum_2Enum^{A_27a}). (\forall V1c \in (ty_2Enum_2Enum^{A_27a}). (((ap\ (ap\ (c_2Ebag_2EBAG_DIFF\ A_27a)\ V0b)\ V1c) = (c_2Ebag_2EEMPTY_BAG\ A_27a)) \Leftrightarrow (p\ (ap\ (ap\ (c_2Ebag_2ESUB_BAG\ A_27a)\ V0b)\ V1c))))))$$