

thm_2Ebag_2EBAG_GEN_SUM_REC (TMZD-PZXWbfKf3ArNWVGc93MW8TrWNkSRYYm)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x)))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 7 We define $c_2Earithmic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmic_2E_2B V0n) V0n)$.

Definition 8 We define $c_2Earithmic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Definition 9 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 10 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 11 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t) V1t2) V0t1))$.

Definition 12 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then (the $(\lambda x.x \in A \wedge P x)$) of type $\iota \Rightarrow \iota$.

Definition 13 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.(ap (c_2Ebool_2E_21 2) (\lambda V2t3 \in 2.V2t3) V2t2) V1t1) V0t))$.

Definition 14 We define $c_2Ebag_2EBAG_INSERT$ to be $\lambda A_27a : \iota.\lambda V0e \in A_27a.\lambda V1b \in (ty_2Enum_2Enum A_27a)$.

Definition 15 We define $c_2Ecombin_2EK$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.(\lambda V0x \in A_27a.(\lambda V1y \in A_27b.V0x))$.

Definition 16 We define $c_2Ebag_2EEMPTY_BAG$ to be $\lambda A_27a : \iota.(ap (c_2Ecombin_2EK ty_2Enum_2Enum A_27a) V0)$.

Definition 17 We define $c_2Ebag_2EFINITE_BAG$ to be $\lambda A_27a : \iota.\lambda V0b \in (ty_2Enum_2Enum^{A_27a}).(ap (c_2Ebag_2EBAG_INSERT ty_2Enum_2Enum^{A_27a} V0b) V0)$.

Let $c_2Ebag_2EITBAG : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Ebag_2EITBAG \\ & A_27a A_27b \in (((A_27b^{A_27b})(ty_2Enum_2Enum^{A_27a}))((A_27b^{A_27b})^{A_27a})) \end{aligned} \quad (7)$$

Definition 18 We define $c_2Ebag_2EBAG_GEN_SUM$ to be $\lambda V0bag \in (ty_2Enum_2Enum^{ty_2Enum_2Enum})$.

Definition 19 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_21 2))$.

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & (ap (ap c_2Earithmic_2E_2B V0m) V1n) = (ap (ap c_2Earithmic_2E_2B \\ & V1n) V0m)))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & \forall V2p \in ty_2Enum_2Enum.((ap (ap c_2Earithmic_2E_2B V0m) \\ & (ap (ap c_2Earithmic_2E_2B V1n) V2p)) = (ap (ap c_2Earithmic_2E_2B \\ & (ap (ap c_2Earithmic_2E_2B V0m) V1n) V2p)))))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned}
& (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (\\
& \quad \forall V2p \in ty_2Enum_2Enum. (((ap (ap c_2Earithmic_2E_2B V0m) \\
& \quad V2p) = (ap (ap c_2Earithmic_2E_2B V1n) V2p)) \Leftrightarrow (V0m = V1n))))))
\end{aligned} \tag{10}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow (\\
& \quad \forall V0f \in ((A_27b^{A_27b})^{A_27a}). (\forall V1e \in A_27a. (\forall V2b \in \\
& \quad (ty_2Enum_2Enum^{A_27a}). (\forall V3a \in A_27b. (((\forall V4x \in A_27a. \\
& \quad (\forall V5y \in A_27a. (\forall V6z \in A_27b. ((ap (ap V0f V4x) (ap (ap \\
& \quad V0f V5y) V6z)) = (ap (ap V0f V5y) (ap (ap V0f V4x) V6z)))))) \wedge (p (ap (\\
& \quad c_2Ebag_2EFINITE_BAG A_27a) V2b))) \Rightarrow ((ap (ap (ap (c_2Ebag_2EITBAG \\
& \quad A_27a A_27b) V0f) (ap (ap (c_2Ebag_2EBAG_INSERT A_27a) V1e) V2b)) \\
& \quad V3a) = (ap (ap V0f V1e) (ap (ap (ap (c_2Ebag_2EITBAG A_27a A_27b) V0f) \\
& \quad V2b) V3a)))))))))
\end{aligned} \tag{11}$$

Assume the following.

$$True \tag{12}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \tag{13}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\
& (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(\\
& \quad p V0t))))))
\end{aligned} \tag{14}$$

Theorem 1

$$\begin{aligned}
& (\forall V0b \in (ty_2Enum_2Enum^{ty_2Enum_2Enum}). ((p (ap (c_2Ebag_2EFINITE_BAG \\
& \quad ty_2Enum_2Enum) V0b)) \Rightarrow (\forall V1x \in ty_2Enum_2Enum. (\forall V2a \in \\
& \quad ty_2Enum_2Enum. ((ap (ap c_2Ebag_2EBAG_GEN_SUM (ap (ap (c_2Ebag_2EBAG_INSERT \\
& \quad ty_2Enum_2Enum) V1x) V0b)) V2a) = (ap (ap c_2Earithmic_2E_2B \\
& \quad V1x) (ap (ap c_2Ebag_2EBAG_GEN_SUM V0b) V2a))))))
\end{aligned}$$