

thm_2Ebag_2EBAG_GEN_SUM_TAILREC
 (TML77DN6LuMxsx5EeTPW4JZLb24oGVmhFTp)

October 26, 2020

Let $c_2Enum_2ZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ZERO_REP \in \omega \quad (1)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$\text{nonempty } ty_2Enum_2Enum \quad (2)$$

Let $c_2Enum_2ABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2ABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (3)$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be ($ap\ c_2Enum_2ABS_num\ c_2Enum_2ZERO_REP$).

Definition 3 We define $c_2Earithmetic_2ZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (4)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (5)$$

Definition 4 We define c_2Ebool_2ET to be ($ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x)$)

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x)))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum. (ap\ c_2Enum_2ABS_num\ (c_2Emin_2E_3D\ (2^{A-27a})))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (6)$$

Definition 7 We define $c_2Earithmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap\ (ap\ c_2Earithmetic_2EBIT1\ n)\ V)$

Definition 8 We define `c_2Earithmetic_2ENUMERAL` to be $\lambda V0x \in ty_2Enum_2Enum. V0x$.

Definition 9 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t\in 2.V0t))$.

Definition 10 We define $c_{\text{2Emin_2E_3D_3D_3E}}$ to be $\lambda P \in 2.\lambda Q \in 2.\text{inj_o} (p \Rightarrow p Q)$ of type ι .

Definition 11 We define $c_{\text{CBool}}(2E_2F_5C)$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap(c_{\text{CBool}}(2E_2F_5C), t1), t2)))$

Definition 12 We define $c_{\text{Emin}}.40$ to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p \text{ (ap } P \text{ } x)) \text{ then } (\lambda x. x \in A \wedge_P x \in P) \text{ else } \perp$ of type $\iota \Rightarrow \iota$.

Definition 13 We define $c_{\text{Ebool_ECOND}}$ to be $\lambda A.27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A.27a.(\lambda V2t2 \in A.27a.($

Definition 14 We define $c_2EBag_EBAG_INSERT$ to be $\lambda A_27a : \iota. \lambda V0e \in A_27a. \lambda V1b \in (ty_2Enum_2E$

Definition 15 We define $c_2Ecombin_2EK$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.(\lambda V0x \in A_27a.(\lambda V1y \in A_27b.V0x))$

Definition 16 We define $c_2Ebag_2EEMPTY_BAG$ to be $\lambda A_27a : \iota.(ap\ (c_2Ecombin_2EK\ ty_2Enum_2En)$

Definition 17 We define $c_2Ebag_EFINITE_BAG$ to be $\lambda A.\lambda 27a : \iota.\lambda V0b \in (ty_2Enum_2Enum^A_\lambda 27a).(ap$

Let $c_{2EBag}_2EITBAG : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$\forall A_27a.\text{nonempty } A_27a \Rightarrow \forall A_27b.\text{nonempty } A_27b \Rightarrow c_2Ebag_2I$

$$A_{\text{27a}} A_{\text{27b}} \in (((A_{\text{27b}}^{A_{\text{27b}}})^{(ty_2E\text{num_2}E\text{num}} \dots))^{((A_{\text{27b}} \dots) \dots)}) \quad (7)$$

Definition 18 We define the **LEADER-AGENT-TERM** to be $\lambda x. \text{seq} \in (\text{seq} \text{--} \text{LEADER-TERM})$

DEFINITION 10 We define CLIQUE-CLIQUE_k to be $(X, \mathcal{C}) \in \Sigma_k \wedge (\exists P \in \text{CLIQUE}_k(\mathcal{C})) \vee (\exists Q \in \text{CLIQUE}_k(\mathcal{C}))$

Assume the following.

$$(ap (ap c_2Earithmetic_2E_2B V0m) V1n) = (ap (ap c_2Earithmetic_2E_2B V1n) V0m))) \quad (8)$$

Assume the following.

Assume the following.

$$\begin{aligned}
 & (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. \\
 & \forall V2p \in ty_2Enum_2Enum. (((ap (ap c_2Earithmetic_2E_2B V0m) \\
 & V2p) = (ap (ap c_2Earithmetic_2E_2B V1n) V2p)) \Leftrightarrow (V0m = V1n)))) \\
 & \tag{10}
 \end{aligned}$$

Assume the following.

$$\begin{aligned}
 & \forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow \\
 & \forall V0f \in ((A_27b^{A_27b})^{A_27a}). (\forall V1b \in (ty_2Enum_2Enum^{A_27a}). \\
 & (((\forall V2x \in A_27a. (\forall V3y \in A_27a. (\forall V4z \in A_27b. \\
 & ((ap (ap V0f V2x) (ap (ap V0f V3y) V4z)) = (ap (ap V0f V3y) (ap (ap V0f \\
 & V2x) V4z))))))) \wedge (p (ap (c_2Ebag_2EFINITE_BAG A_27a) V1b))) \Rightarrow \\
 & \forall V5x \in A_27a. (\forall V6a \in A_27b. ((ap (ap (ap (c_2Ebag_2EITBAG \\
 & A_27a A_27b) V0f) (ap (ap (c_2Ebag_2EBAG_INSERT A_27a) V5x) V1b)) \\
 & V6a) = (ap (ap (ap (c_2Ebag_2EITBAG A_27a A_27b) V0f) V1b) (ap (ap \\
 & V0f V5x) V6a))))))) \\
 & \tag{11}
 \end{aligned}$$

Assume the following.

$$True \tag{12}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \tag{13}$$

Assume the following.

$$\begin{aligned}
 & (\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\
 & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p \\
 & V0t))))))) \\
 & \tag{14}
 \end{aligned}$$

Theorem 1

$$\begin{aligned}
 & (\forall V0b \in (ty_2Enum_2Enum^{ty_2Enum_2Enum}). ((p (ap (c_2Ebag_2EFINITE_BAG \\
 & ty_2Enum_2Enum) V0b)) \Rightarrow (\forall V1x \in ty_2Enum_2Enum. (\forall V2a \in \\
 & ty_2Enum_2Enum. ((ap (ap c_2Ebag_2EBAG_GEN_SUM (ap (ap (c_2Ebag_2EBAG_INSERT \\
 & ty_2Enum_2Enum) V1x) V0b)) V2a) = (ap (ap c_2Ebag_2EBAG_GEN_SUM \\
 & V0b) (ap (ap c_2Earithmetic_2E_2B V1x) V2a)))))))
 \end{aligned}$$