

thm\_2Ebag\_2EBAG\_REST\_SING  
(TMc2dtZWC1d6sm2zuoM2JGNgZHezZ9RSHjc)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow p Q)$  of type  $\iota$ .

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P)))$

**Definition 5** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))))$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (1)$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (2)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (3)$$

**Definition 6** We define  $c\_2Enum\_2E0$  to be  $(ap c\_2Enum\_2EABS\_num c\_2Enum\_2EZERO\_REP)$ .

**Definition 7** We define  $c\_2Ecombin\_2EK$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.(\lambda V0x \in A\_27a.(\lambda V1y \in A\_27b.V0x))$

**Definition 8** We define  $c\_2Ebag\_2EEMPTY\_BAG$  to be  $\lambda A\_27a : \iota.(ap (c\_2Ecombin\_2EK ty\_2Enum\_2Enum) (c\_2Eempty\_2BAG))$

**Definition 9** We define  $c\_2Earithmetic\_2EZERO$  to be  $c\_2Enum\_2E0$ .

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \quad (4)$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \quad (5)$$

**Definition 10** We define  $c_2Enum_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c_2Enum_2EABS\_num$   
 Let  $c_2Earithmetic_2E_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^*ty\_2Enum\_2Enum)^*)ty\_2Enum\_2Enum \quad (6)$$

**Definition 11** We define  $c\_2Earithmetic\_2EBIT1$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap\ (ap\ c\_2Earithmetic\ 0\ n)\ V)$

**Definition 12** We define `c_2Earthmetic_2ENUMERAL` to be  $\lambda V0x \in ty\_2Enum\_2Enum. V0x$ .

**Definition 13** We define  $c\_2Ebool\_2EF$  to be  $(ap\ (c\_2Ebool\_2E_21\ 2)\ (\lambda V0t\ t \in 2.V0t))$ .

**Definition 14** We define  $c_2 \in \text{min}_2 \in \text{A} \cdot \text{if } (\exists x \in A.p \text{ (ap } P \text{ } x)) \text{ then } P \text{ else } Q$

**Definition 15** We define  $\text{c\_2Epool\_2ECOND}$  to be  $\lambda A.\lambda 27a:\iota_1.(\lambda V0t \in 2.(\lambda V1t1 \in A.27a.(\lambda V2t2 \in A.27a.($

**Definition 16** We define  $c \in \text{Ebag} \cap \text{EBAG}$ . **INSERT** to be  $\lambda A. 2\pi a : t \cdot \lambda V0e \in A. 2\pi a \cdot \lambda V1b \in (tu : \text{Enum} \cap \text{Ebag})$ .

**Definition 17.** We define a 2Ebag-2EEFL<sub>1</sub>-PAC to be  $\lambda(4, 27c) \cup \lambda(V_0c \in 4, 27c)$  (or  $\lambda(c \in 4, 27c)$ ) if  $c$  is a 2Ebag-2EEFL<sub>1</sub>-PAC.

Let  $\Delta E$  with unit  $\Delta E$  2D and  $\alpha = \Delta E$  with the following:

<sup>2</sup>  $\Sigma_1 = \{1\}$ ,  $\Sigma_2 = \{1, 2\}$ ,  $\Sigma_3 = \{1, 2, 3\}$ ,  $\Sigma_4 = \{1, 2, 3, 4\}$ ,  $\Sigma_5 = \{1, 2, 3, 4, 5\}$ ,  $\Sigma_6 = \{1, 2, 3, 4, 5, 6\}$ ,  $\Sigma_7 = \{1, 2, 3, 4, 5, 6, 7\}$ ,  $\Sigma_8 = \{1, 2, 3, 4, 5, 6, 7, 8\}$ .

$$e_2 \text{Barinmetric}_2 E_2 D \in ((g_2 \text{Enam}_2 \text{Enam})^*)^* \quad (7)$$

**Definition 18** We define  $\text{C2EBag-ZEBAG-DIFF}$  to be  $\text{XA-ZTA} : \lambda x. \text{box} \in (\text{ty-ZENAM-ZENAM}) \rightarrow \lambda y. \text{box}$

Let  $c\_2Day\_2EBAG\_CHOICE : t \rightarrow t$  be given. Assume the following.

$$\forall A \exists Z (A \text{ nonempty} \wedge A \subseteq Z \wedge \forall a \in A \exists x \in Z \forall y \in Z (x \in y \rightarrow a \in y)) \quad (8)$$

**Definition 19** We define  $c\_2EBag\_2EBAG\_REST$  to be  $\lambda A\_\underline{27}a : \iota.\lambda V0b \in (ty\_2Enum\_2Enum^{\wedge-27a}).(ap (d$

Assume the following.

$$\begin{aligned} & \forall A\_27a.\text{nonempty } A\_27a \Rightarrow \forall A\_27b.\text{nonempty } A\_27b \Rightarrow \forall A\_27c. \\ & \quad \text{nonempty } A\_27c \Rightarrow ((\forall V0b \in (\text{ty\_2Enum\_2Enum}^{A\_27a})).((ap \ (ap \\ & \quad (c\_2Ebag\_2EBAG\_DIFF \ A\_27a) \ V0b) \ V0b) = (c\_2Ebag\_2EEMPTY\_BAG \\ & \quad A\_27a))) \wedge ((\forall V1b \in (\text{ty\_2Enum\_2Enum}^{A\_27b})).((ap \ (ap \ (c\_2Ebag\_2EBAG\_DIFF \\ & \quad A\_27b) \ V1b) \ (c\_2Ebag\_2EEMPTY\_BAG \ A\_27b)) = V1b)) \wedge (\forall V2b \in \\ & \quad (\text{ty\_2Enum\_2Enum}^{A\_27c})).((ap \ (ap \ (c\_2Ebag\_2EBAG\_DIFF \ A\_27c) \\ & \quad (c\_2Ebag\_2EEMPTY\_BAG \ A\_27c)) \ V2b) = (c\_2Ebag\_2EEMPTY\_BAG \ A\_27c)))))) \\ & \quad (9) \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.\text{nonempty } A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1b1 \in \\ & (ty\_2Enum\_2Enum^{A\_27a}).(\forall V2b2 \in (ty\_2Enum\_2Enum^{A\_27a}). \\ & ((ap (ap (c\_2Ebag\_2EBAG\_DIFF A\_27a) (ap (ap (c\_2Ebag\_2EBAG\_INSERT \\ & A\_27a) V0x) V1b1)) (ap (ap (c\_2Ebag\_2EBAG\_INSERT A\_27a) V0x) V2b2)) = \\ & (ap (ap (c\_2Ebag\_2EBAG\_DIFF A\_27a) V1b1) V2b2)))))) \\ & \end{aligned} \quad (10)$$

Assume the following.

$$\forall A_27a.\text{nonempty } A_27a \Rightarrow (\forall V0x \in A_27a. ((\text{ap } (\text{c\_2EBAG\_2EBAG\_CHOICE } A_27a) (\text{ap } (\text{ap } (\text{c\_2EBAG\_2EBAG\_INSERT } A_27a) V0x) (\text{c\_2EBAG\_2EEMPTY\_BAG } A_27a))) = V0x)) \quad (11)$$

Assume the following.

$$True \quad (12)$$

Assume the following.

$$\forall A_27a.\text{nonempty } A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (13)$$

### Theorem 1

$$\forall A_27a.\text{nonempty } A_27a \Rightarrow (\forall V0x \in A_27a. ((\text{ap } (\text{c\_2EBAG\_2EBAG\_REST } A_27a) (\text{ap } (\text{ap } (\text{c\_2EBAG\_2EBAG\_INSERT } A_27a) V0x) (\text{c\_2EBAG\_2EEMPTY\_BAG } A_27a))) = (\text{c\_2EBAG\_2EEMPTY\_BAG } A_27a)))$$