

thm_2Ebag_2EBAG__UNION__DIFF__eliminate (TMW7z9cb26T4oCF5Z93Qzvyuq8PjWVTMEwq)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow P \Rightarrow Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 5 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{2}$$

Definition 6 We define $c_2Ebag_2EBAG_UNION$ to be $\lambda A_27a : \iota.\lambda V0b \in (ty_2Enum_2Enum^{A-27a}).\lambda V1c \in (ty_2Enum_2Enum^{A-27a}).$

Let $c_2Earithmetic_2E_2D : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2D \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{3}$$

Definition 7 We define $c_2Ebag_2EBAG_DIFF$ to be $\lambda A_27a : \iota.\lambda V0b1 \in (ty_2Enum_2Enum^{A-27a}).\lambda V1b2 \in (ty_2Enum_2Enum^{A-27a}).$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & (ap (ap c_2Earithmetic_2E_2B V0m) V1n) = (ap (ap c_2Earithmetic_2E_2B \\ & V1n) V0m)))) \end{aligned} \tag{4}$$

Assume the following.

$$(\forall V0a \in ty_2Enum_2Enum. (\forall V1c \in ty_2Enum_2Enum. (ap (ap c_2Earithmic_2E_2D (ap (ap c_2Earithmic_2E_2B V0a) V1c)) V1c) = V0a))) \quad (5)$$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (7)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (8)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow (\forall V0f \in (A_27b^{A_27a}). (\forall V1g \in (A_27b^{A_27a}). ((V0f = V1g) \Leftrightarrow (\forall V2x \in A_27a. ((ap V0f V2x) = (ap V1g V2x)))))) \quad (9)$$

Theorem 1

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0b \in (ty_2Enum_2Enum^{A_27a}). (\forall V1c \in (ty_2Enum_2Enum^{A_27a}). (((ap (ap (c_2Ebag_2EBAG_DIFF A_27a) (ap (ap (c_2Ebag_2EBAG_UNION A_27a) V0b) V1c)) V1c) = V0b) \wedge ((ap (ap (c_2Ebag_2EBAG_DIFF A_27a) (ap (ap (c_2Ebag_2EBAG_UNION A_27a) V1c) V0b)) V1c) = V0b))))))$$