

thm_2Ebag_2EFINITE_BAG_INDUCT
 (TMThEXH-
 Jag45EZ7eUTHS2TwU1yQLmdiGAeY)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (1)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$\text{nonempty } ty_2Enum_2Enum \quad (2)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (3)$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (4)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (5)$$

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ (ap\ (c_2Emin_2E_3D\ (2^{A_27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x)))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum. (ap\ c_2Enum_2EABS_num\ m)$

Let c_2 be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (6)$$

Definition 7 We define $c_2Earithmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap\ (ap\ c_2Earithmetic_2EBIT1\ n)\ V)$

Definition 8 We define c_2Earithmetic_2ENUMERAL to be $\lambda V0x \in ty_2Enum_2Enum. V0x.$

Definition 9 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t\ t \in 2.V0t))$.

Definition 10 We define $c_2 E_{\min} 2E_3 D_3 D_3 E$ to be $\lambda P \in 2. \lambda Q \in 2. inj_o (p \ P \Rightarrow p)$

Definition 11 We define $c_2\text{Ebool}$, $c_2\text{E}$, $c_2\text{C}$, $c_2\text{S}$ to be $(\lambda V0t1 \in 2)(\lambda V1t2 \in 2)$ (α) $(c_2\text{Ebool})$

Definition 12. We define \subseteq 2Emin 2E 40 to be $\lambda A.\lambda B\in 2^A$ if $(\exists x\in A.p.(ap.B.x))$, then (the $\lambda x.x\in A$)

of type $\iota \mapsto \iota$.

Definition 10 We define **COLLECTIVE** to be $\text{AFL}(\text{COLL}_\infty \cup \{\lambda\})$, where $\text{COLL}_\infty \subseteq \text{AFL}(\text{COLL}_1)$, $\text{COLL}_1 \subseteq \text{AFL}(\text{COLL}_0)$, $\text{COLL}_0 \subseteq \text{AFL}(\text{COLL}_\infty)$.

Definition 18 We define $c_{\text{ZEBAG-ZELEM}}: \text{ZEBAG} \rightarrow \text{ZELEM}$ to be $\lambda A \lambda x. a : i. (ap (c_{\text{ZEBAG-CELEM}}) (g_{\text{ZEBAG-CELEM}}))$

Definition 17 We define $c_2Ebag_2EFINITE_BAG$ to be $\lambda A.\exists a : \iota. \lambda V. 0b \in (ty_enum_enum^{\iota \rightarrow \tau_a}).(ap$

Definition 18 We define $c_2Eb0l_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Eb0l_2E_7E))$

Assume the following.

True (7)

Assume the following.

$$\forall A_27a.\text{nonempty } A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (8)$$

Assume the following.

$$(\forall V0t \in 2.(((True \wedge (p \vee 0t)) \Leftrightarrow (p \vee 0t)) \wedge (((p \vee 0t) \wedge True) \Leftrightarrow (p \vee 0t)) \wedge (((False \wedge (p \vee 0t)) \Leftrightarrow False) \wedge (((p \vee 0t) \wedge False) \Leftrightarrow False) \wedge (((p \vee 0t) \wedge (p \vee 0t)) \Leftrightarrow (p \vee 0t))))))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p \ V0t)) \Leftrightarrow True) \wedge (((p \ V0t) \Rightarrow (p \ V0t)) \Leftrightarrow True) \wedge (((p \ V0t) \Rightarrow False) \Leftrightarrow (\neg(p \ V0t))))))) \quad (10)$$

Assume the following.

$$\forall A_{\text{27a}}. \text{nonempty } A_{\text{27a}} \Rightarrow (\forall V0x \in A_{\text{27a}}. (\forall V1y \in A_{\text{27a}}. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (11)$$

Assume the following.

$$(\forall V0t \in 2. (((\text{True} \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow \text{True}) \Leftrightarrow (p V0t))) \wedge (((\text{False} \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow \text{False}) \Leftrightarrow (\neg(p V0t))))) \quad (12)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3))))) \quad (13)$$

Assume the following.

$$(\forall V0x \in 2. (\forall V1x_{\text{27}} \in 2. (\forall V2y \in 2. (\forall V3y_{\text{27}} \in 2. (((p V0x) \Leftrightarrow (p V1x_{\text{27}})) \wedge ((p V1x_{\text{27}}) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_{\text{27}})))))) \Rightarrow ((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_{\text{27}}) \Rightarrow (p V3y_{\text{27}})))) \quad (14)$$

Theorem 1

$$\begin{aligned} & \forall A_{\text{27a}}. \text{nonempty } A_{\text{27a}} \Rightarrow (\forall V0P \in (2^{(ty_{\text{2Enum}} \cdot 2^{Enum^{A_{\text{27a}}}})}). \\ & (((p (ap V0P (c_{\text{2EBAG_2EMPTY_BAG}} A_{\text{27a}}))) \wedge (\forall V1b \in (ty_{\text{2Enum}} \cdot 2^{Enum^{A_{\text{27a}}}}). \\ & ((p (ap V0P V1b)) \Rightarrow (\forall V2e \in A_{\text{27a}}. (p (ap V0P (ap (ap (c_{\text{2EBAG_2EBAG_INSERT}} \\ & A_{\text{27a}}) V2e) V1b))))))) \Rightarrow (\forall V3b \in (ty_{\text{2Enum}} \cdot 2^{Enum^{A_{\text{27a}}}}). \\ & ((p (ap (c_{\text{2EBAG_2FINITE_BAG}} A_{\text{27a}}) V3b)) \Rightarrow (p (ap V0P V3b))))))) \end{aligned}$$