

thm_2Ebag_2EFINITE__EMPTY__BAG (TM- byF6zdGEWJUR2hFYm1wqgeMLWzKNDDWHw)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 7 We define `c_2Earithmic_2EBIT1` to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmic_2EBIT1) V0n)$.

Definition 8 We define `c_2Earithmic_2ENUMERAL` to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Definition 9 We define `c_2Ebool_2EF` to be $(ap (c_2Ebool_2E_21) 2) (\lambda V0t \in 2.V0t)$.

Definition 10 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 11 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21) 2) (\lambda V2t \in 2.V2t) V0t1 V1t2))$.

Definition 12 We define `c_2Emin_2E_40` to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then $(the (\lambda x.x \in A \wedge P x))$ of type $\iota \Rightarrow \iota$.

Definition 13 We define `c_2Ebool_2ECOND` to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.(ap (c_2Ebool_2E_21) 2) V0t1 V2t2) V1t1))$.

Definition 14 We define `c_2Ebag_2EBAG_INSERT` to be $\lambda A_27a : \iota.\lambda V0e \in A_27a.\lambda V1b \in (ty_2Enum_2Enum A_27a).(\lambda V2c \in A_27a.(ap (c_2Ebag_2EBAG_INSERT) V0e V1b) V2c))$.

Definition 15 We define `c_2Ecombin_2EK` to be $\lambda A_27a : \iota.\lambda A_27b : \iota.(\lambda V0x \in A_27a.(\lambda V1y \in A_27b.V0x))$.

Definition 16 We define `c_2Ebag_2EEMPTY_BAG` to be $\lambda A_27a : \iota.(ap (c_2Ecombin_2EK) ty_2Enum_2Enum A_27a)$.

Definition 17 We define `c_2Ebag_2EFINITE_BAG` to be $\lambda A_27a : \iota.\lambda V0b \in (ty_2Enum_2Enum^{A_27a}).(ap (c_2Ebag_2EBAG_INSERT) V0b)$.

Definition 18 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E) V0t) c_2Ebool_2E_7E)$.

Assume the following.

$$True \tag{7}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \tag{8}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \end{aligned} \tag{9}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{10}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg (p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg \\ & (p V0t)))))) \end{aligned} \tag{11}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (12)$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1x_{.27} \in 2.(\forall V2y \in 2.(\forall V3y_{.27} \in 2.(((p V0x) \Leftrightarrow (p V1x_{.27})) \wedge ((p V1x_{.27}) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_{.27})))))) \Rightarrow (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_{.27}) \Rightarrow (p V3y_{.27})))))) \quad (13)$$

Theorem 1

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow (p (ap (c_{.2Ebag_{.2EFINITE_BAG} A_{.27a}} (c_{.2Ebag_{.2EEMPTY_BAG} A_{.27a}})))$$