

# thm\_2Ebag\_2ENOT\_\_mlt\_\_EMPTY (TMYFh- LLAdCSJR2EZvv2gHarTy6zhECAvNiG)

October 26, 2020

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{1}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{2}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{3}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

**Definition 3** We define  $c\_2Ecombin\_2EK$  to be  $\lambda A.\lambda a : \iota.\lambda A.\lambda b : \iota.(\lambda V0x \in A.\lambda V1y \in A.V0x)$

**Definition 4** We define  $c\_2Ebag\_2EEMPTY\_BAG$  to be  $\lambda A.\lambda a : \iota.(ap\ (c\_2Ecombin\_2EK\ ty\_2Enum\_2Enum))$

**Definition 5** We define  $c\_2Earithmic\_2EZERO$  to be  $c\_2Enum\_2E0$ .

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{4}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{5}$$

**Definition 6** We define  $c\_2Ebool\_2ET$  to be  $(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 7** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^{A-27a}))))$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num\ (c\_2Ecombin\_2EK\ ty\_2Enum\_2Enum))$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})ty\_2Enum\_2Enum) \quad (6)$$

**Definition 9** We define  $c\_2Earithmetic\_2EBIT1$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap (ap c\_2Earithmetic\_2E\_2B) V0n)$ .

**Definition 10** We define  $c\_2Earithmetic\_2ENUMERAL$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

**Definition 11** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21) 2) (\lambda V0t \in 2.V0t)$ .

**Definition 12** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 13** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E) V0t) c\_2Ebool\_2E\_21)$ .

**Definition 14** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21) 2) (\lambda V2t \in 2.V2t)))$ .

**Definition 15** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$  then  $(the (\lambda x.x \in A \wedge P x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 16** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40) V0P)))$ .

**Definition 17** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.V1n$ .

**Definition 18** We define  $c\_2Earithmetic\_2E\_3E$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.V1n$ .

**Definition 19** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21) 2) (\lambda V2t \in 2.V2t)))$ .

**Definition 20** We define  $c\_2Earithmetic\_2E\_3E\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.V1n$ .

**Definition 21** We define  $c\_2Ebag\_2EBAG\_INN$  to be  $\lambda A\_27a : \iota.\lambda V0e \in A\_27a.\lambda V1n \in ty\_2Enum\_2Enum.V1n$ .

**Definition 22** We define  $c\_2Ebag\_2EBAG\_IN$  to be  $\lambda A\_27a : \iota.\lambda V0e \in A\_27a.\lambda V1b \in (ty\_2Enum\_2Enum^{A\_27a}).V1b$ .

**Definition 23** We define  $c\_2Ebool\_2ECOND$  to be  $\lambda A\_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A\_27a.(\lambda V2t2 \in A\_27a.V2t2)))$ .

**Definition 24** We define  $c\_2Ebag\_2EBAG\_INSERT$  to be  $\lambda A\_27a : \iota.\lambda V0e \in A\_27a.\lambda V1b \in (ty\_2Enum\_2Enum^{A\_27a}).V1b$ .

**Definition 25** We define  $c\_2Ebag\_2EBAG\_UNION$  to be  $\lambda A\_27a : \iota.\lambda V0b \in (ty\_2Enum\_2Enum^{A\_27a}).V0b$ .

**Definition 26** We define  $c\_2Ebag\_2EFINITE\_BAG$  to be  $\lambda A\_27a : \iota.\lambda V0b \in (ty\_2Enum\_2Enum^{A\_27a}).(ap V0b)$ .

**Definition 27** We define  $c\_2Ebag\_2Emlt1$  to be  $\lambda A\_27a : \iota.\lambda V0r \in ((2^{A\_27a})^{A\_27a}).\lambda V1b1 \in (ty\_2Enum\_2Enum^{A\_27a}).V1b1$ .

**Definition 28** We define  $c\_2Ebool\_2EBOUNDED$  to be  $(\lambda V0v \in 2.c\_2Ebool\_2ET)$ .

**Definition 29** We define  $c\_2ERelation\_2ETC$  to be  $\lambda A\_27a : \iota.\lambda V0R \in ((2^{A\_27a})^{A\_27a}).\lambda V1a \in A\_27a.\lambda V2b \in A\_27a.V2b$ .

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0r \in ((2^{A\_27a})^{A\_27a}). \\ & (\forall V1b \in (ty\_2Enum\_2Enum^{A\_27a}). (\neg(p\ (ap\ (ap\ (ap\ (c\_2Ebag\_2Emlt1 \\ & A\_27a)\ V0r)\ V1b)\ (c\_2Ebag\_2EEMPTY\_BAG\ A\_27a)))))) \end{aligned} \quad (7)$$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p\ V0t))) \quad (10)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0t \in 2. ((\exists V1x \in A\_27a. (p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (11)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (12)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge \\ & (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee \\ & (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (13)$$

Assume the following.

$$((\forall V0t \in 2. ((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t)) \wedge ((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (14)$$

Assume the following.

$$(\forall V0v \in 2. ((p\ (ap\ c\_2Ebool\_2EBOUNDED\ V0v)) \Leftrightarrow True)) \quad (15)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0R \in ((2^{A\_27a})^{A\_27a}). \\ & (\forall V1x \in A\_27a. (\forall V2z \in A\_27a. ((p\ (ap\ (ap\ (ap\ (c\_2Erelation\_2ETC \\ & A\_27a)\ V0R)\ V1x)\ V2z)) \Leftrightarrow ((p\ (ap\ (ap\ V0R\ V1x)\ V2z)) \vee (\exists V3y \in A\_27a. \\ & ((p\ (ap\ (ap\ (ap\ (c\_2Erelation\_2ETC\ A\_27a)\ V0R)\ V1x)\ V3y)) \wedge (p\ (ap \\ & (ap\ V0R\ V3y)\ V2z)))))))))) \end{aligned} \quad (16)$$

**Theorem 1**

$$\begin{aligned} & \forall A_{27a}. \text{nonempty } A_{27a} \Rightarrow (\forall V0R \in ((2^{A_{27a}})^{A_{27a}}). \\ & (\forall V1b \in (\text{ty\_2Enum\_2Enum}^{A_{27a}}). (\neg (p \text{ (ap (ap (ap (c\_2Erelation\_2ETC} \\ & (\text{ty\_2Enum\_2Enum}^{A_{27a}})) \text{ (ap (c\_2Ebag\_2Emlt1 } A_{27a}) V0R)) V1b) \\ & (\text{c\_2Ebag\_2EMPTY\_BAG } A_{27a})))))) \end{aligned}$$