

thm_2Ebag_2EPSUB__BAG__NOT__EQ (TMQbVG6SBipde9LhYfQrfJF9kiXF1VnTzLH)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num (ap c_2Enum_2EREP_num (ap c_2Enum_2ESUC_REP m)))$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge P x))$ of type $\iota \Rightarrow \iota$.

Definition 10 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ V0P\ (ap\ (c_2Emin_2E_40$

Definition 11 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Definition 12 We define $c_2Earithmetic_2E_3E$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Definition 13 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in$

Definition 14 We define $c_2Earithmetic_2E_3E_3D$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Definition 15 We define $c_2Ebag_2EBAG_INN$ to be $\lambda A_27a : \iota. \lambda V0e \in A_27a. \lambda V1n \in ty_2Enum_2Enum$

Definition 16 We define $c_2Ebag_2ESUB_BAG$ to be $\lambda A_27a : \iota. \lambda V0b1 \in (ty_2Enum_2Enum^{A_27a}). \lambda V1b2 \in$

Definition 17 We define $c_2Ebag_2EPSUB_BAG$ to be $\lambda A_27a : \iota. \lambda V0b1 \in (ty_2Enum_2Enum^{A_27a}). \lambda V1b2 \in$

Assume the following.

$$True \tag{5}$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \tag{6}$$

Assume the following.

$$\forall A_27a. nonempty\ A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_27a. (p\ V0t)) \Leftrightarrow (p\ V0t))) \tag{7}$$

Assume the following.

$$(\forall V0t \in 2. (((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \tag{8}$$

Assume the following.

$$((\forall V0t \in 2. ((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \tag{9}$$

Assume the following.

$$\forall A_27a. nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \tag{10}$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \tag{11}$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((p V0A) \vee (p V1B)) \Leftrightarrow ((p V1B) \vee (p V0A)))) \quad (12)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((p V0A) \Rightarrow (p V1B)) \Leftrightarrow ((\neg(p V0A)) \vee (p V1B)))) \quad (13)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0f \in (2^{A_27a}).(\forall V1v \in A_27a.((\forall V2x \in A_27a.((V2x = V1v) \Rightarrow (p (ap V0f V2x)))) \Leftrightarrow (p (ap V0f V1v)))))) \quad (14)$$

Theorem 1

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0b1 \in (ty_2Enum_2Enum^{A_27a}).(\forall V1b2 \in (ty_2Enum_2Enum^{A_27a}).((p (ap (ap (c_2Ebag_2EPSUB_BAG A_27a) V0b1) V1b2)) \Rightarrow (\neg(V0b1 = V1b2))))))$$