

thm_2Ebag_2ESING__EL__BAG
(TMY8UDn4QNNQeCypXP5tRabCphqQnfYqwumt)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Ecombin_2EK$ to be $\lambda A.\lambda a : \iota.\lambda A.\lambda b : \iota.(\lambda V0x \in A.\lambda V1y \in A.V0x)$

Definition 4 We define $c_2Ebag_2EEMPTY_BAG$ to be $\lambda A.\lambda a : \iota.(ap\ (c_2Ecombin_2EK\ ty_2Enum_2Enum))$

Definition 5 We define $c_2Earithmic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 6 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 7 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))))$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (c_2Ecombin_2EK\ ty_2Enum_2Enum))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (6)$$

Definition 9 We define $c_2Earithmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmetic_2E_2B))$

Definition 10 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Definition 11 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2)) (\lambda V0t \in 2.V0t)$.

Definition 12 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 13 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2)) (\lambda V2t \in 2.V2t)))$

Definition 14 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then $(the (\lambda x.x \in A \wedge P x))$ of type $\iota \Rightarrow \iota$.

Definition 15 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.V2t2)))$

Definition 16 We define $c_2Ebag_2EBAG_INSERT$ to be $\lambda A_27a : \iota.\lambda V0e \in A_27a.\lambda V1b \in (ty_2Enum_2Enum^{A_27a})$

Definition 17 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40))))$

Definition 18 We define $c_2Ebag_2ESING_BAG$ to be $\lambda A_27a : \iota.\lambda V0b \in (ty_2Enum_2Enum^{A_27a}).(ap (c_2Emin_2E_40))$

Definition 19 We define $c_2Ebag_2EEL_BAG$ to be $\lambda A_27a : \iota.\lambda V0e \in A_27a.(ap (ap (c_2Ebag_2EBAG_INSERT)))$

Definition 20 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_21))$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0e \in A_27a.(p (ap (c_2Ebag_2ESING_BAG A_27a) (ap (ap (c_2Ebag_2EBAG_INSERT A_27a) V0e) (c_2Ebag_2EEMPTY_BAG A_27a)))))) \quad (7)$$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg (p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg (p V0t))))) \quad (10)$$

Theorem 1

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0e \in A_27a.(p (ap (c_2Ebag_2ESING_BAG A_27a) (ap (c_2Ebag_2EEL_BAG A_27a) V0e))))$$