

thm\_2Ebit\_2EBITS\_\_DIV\_\_THM (TMY-oVTQesVdgNiWDWgHRaJ2F6TpI7h2K5Ph)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (1)$$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (2)$$

Let  $c\_2Earithmetic\_2E\_2A : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2A \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (3)$$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (4)$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (5)$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A. \lambda x \in A. \lambda y \in A. inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

**Definition 3** We define  $c\_2Ebool\_2ET$  to be  $(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^{A\_27a}))\ (\lambda V1t \in 2.V1t))\ P)))$

**Definition 5** We define  $c\_2Ebool\_2EF$  to be  $(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V0t \in 2.V0t))$ .

**Definition 6** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2. \lambda Q \in 2. inj\_o (p\ P \Rightarrow p\ Q)$  of type  $\iota$ .

**Definition 7** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2. (ap\ (ap\ c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ c\_2Ebool\_2EF))$

**Definition 8** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap(c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \quad (6)$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \quad (7)$$

**Definition 9** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num ($

**Definition 10** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap P x)) \text{ then } (\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ .

**Definition 11** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40$

**Definition 12** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 13** We define  $c\_2Earithmetic\_2EZERO$  to be  $c\_2Enum\_2E0.$

**Definition 14** We define  $c\_2Earithmetic\_2EBIT2$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap (ap c\_2Earithmetic\_2EZERO$

**Definition 15** We define  $c\_2Earithmetic\_2ENUMERAL$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x.$

Let  $c\_2Earithmetic\_2EEEXP : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2EEEXP \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (8)$$

Let  $c\_2Earithmetic\_2EMOD : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2EMOD \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (9)$$

Let  $c\_2Earithmetic\_2EDIV : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2EDIV \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (10)$$

**Definition 16** We define  $c\_2Ebit\_2EDIV\_2EXP$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.$

Let  $c\_2Earithmetic\_2E\_2D : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2D \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (11)$$

**Definition 17** We define  $c\_2Ebit\_2EMOD\_2EXP$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 18** We define  $c\_2Ebit\_2EBITS$  to be  $\lambda V0h \in ty\_2Enum\_2Enum.\lambda V1l \in ty\_2Enum\_2Enum.\lambda V2t \in 2.$

Assume the following.

$$\begin{aligned}
 & (\forall V0p \in ty\_2Enum\_2Enum. (\forall V1q \in ty\_2Enum\_2Enum. (\forall V2n \in ty\_2Enum\_2Enum. ((ap (ap c\_2Earithmetic\_2EEEXP V2n) \\
 & (ap (ap c\_2Earithmetic\_2E\_2B V0p) V1q)) = (ap (ap c\_2Earithmetic\_2E\_2A \\
 & (ap (ap c\_2Earithmetic\_2EEEXP V2n) V0p)) (ap (ap c\_2Earithmetic\_2EEEXP \\
 & V2n) V1q)))))))
 \end{aligned} \tag{12}$$

Assume the following.

$$\begin{aligned}
 & (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. ((p (ap (ap c\_2Eprim\_rec\_2E\_3C c\_2Enum\_2E0) V0m) \wedge (p (ap (ap c\_2Eprim\_rec\_2E\_3C \\
 & c\_2Enum\_2E0) V1n)) \Rightarrow (\forall V2x \in ty\_2Enum\_2Enum. ((ap (ap c\_2Earithmetic\_2EDIV \\
 & (ap (ap c\_2Earithmetic\_2EDIV V2x) V0m)) V1n) = (ap (ap c\_2Earithmetic\_2EDIV \\
 & V2x) (ap (ap c\_2Earithmetic\_2E\_2A V0m) V1n)))))))
 \end{aligned} \tag{13}$$

Assume the following.

$$\begin{aligned}
 & (\forall V0n \in ty\_2Enum\_2Enum. (p (ap (ap c\_2Eprim\_rec\_2E\_3C c\_2Enum\_2E0) \\
 & (ap (ap c\_2Earithmetic\_2EEEXP (ap c\_2Earithmetic\_2ENUMERAL (ap \\
 & c\_2Earithmetic\_2EBIT2 c\_2Earithmetic\_2EZERO)) V0n))))))
 \end{aligned} \tag{14}$$

Assume the following.

$$\begin{aligned}
 & (\forall V0h \in ty\_2Enum\_2Enum. (\forall V1l \in ty\_2Enum\_2Enum. (\forall V2n \in ty\_2Enum\_2Enum. ((ap (ap (ap c\_2Ebit\_2EBITS V0h) V1l) \\
 & V2n) = (ap (ap c\_2Earithmetic\_2EDIV (ap (ap c\_2Earithmetic\_2EMOD \\
 & V2n) (ap (ap c\_2Earithmetic\_2EEEXP (ap c\_2Earithmetic\_2ENUMERAL \\
 & (ap c\_2Earithmetic\_2EBIT2 c\_2Earithmetic\_2EZERO)) (ap c\_2Enum\_2ESUC \\
 & V0h))) (ap (ap c\_2Earithmetic\_2EEEXP (ap c\_2Earithmetic\_2ENUMERAL \\
 & (ap c\_2Earithmetic\_2EBIT2 c\_2Earithmetic\_2EZERO)) V1l)))))))
 \end{aligned} \tag{15}$$

Assume the following.

$$True \tag{16}$$

Assume the following.

$$\begin{aligned}
 & (\forall V0t \in 2. (((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\
 & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\
 & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))))
 \end{aligned} \tag{17}$$

Assume the following.

$$\forall A\_27a.\text{nonempty } A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \tag{18}$$

Assume the following.

$$\begin{aligned}
 & (\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\
 & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg \\
 & p V0t)))))))
 \end{aligned} \tag{19}$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (20)$$

**Theorem 1**

$$\begin{aligned} & (\forall V0h \in ty\_2Enum\_2Enum. (\forall V1l \in ty\_2Enum\_2Enum. ( \\ & \quad \forall V2x \in ty\_2Enum\_2Enum. (\forall V3n \in ty\_2Enum\_2Enum. (( \\ & \quad \quad ap (ap c\_2Earithmetic\_2EDIV (ap (ap (ap c\_2Ebit\_2EBITS V0h) V1l) \\ & \quad \quad V2x)) (ap (ap c\_2Earithmetic\_2EXP (ap c\_2Earithmetic\_2ENUMERAL \\ & \quad \quad (ap c\_2Earithmetic\_2EBIT2 c\_2Earithmetic\_2EZERO))) V3n)) = ( \\ & \quad \quad ap (ap (ap c\_2Ebit\_2EBITS V0h) (ap (ap c\_2Earithmetic\_2E\_2B V1l) \\ & \quad \quad V3n)) V2x))))))) \end{aligned}$$