

thm\_2Ebit\_2EDIV\_\_MULT\_\_1  
(TMGsKvT6sKacTKyQsBnBt6bSHBkanDLLRvy)

October 26, 2020

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{1}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{2}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{3}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

**Definition 3** We define  $c\_2Earithmetic\_2EZERO$  to be  $c\_2Enum\_2E0$ .

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{4}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{5}$$

**Definition 4** We define  $c\_2Ebool\_2ET$  to be  $(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 5** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x)))$

**Definition 6** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num\ (c\_2ESUC\_REP\ m))$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{6}$$

**Definition 7** We define  $c\_2Earithmic\_2EBIT1$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap (ap c\_2Earithmic\_2E\_2A$

**Definition 8** We define  $c\_2Earithmic\_2ENUMERAL$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

Let  $c\_2Earithmic\_2E\_2A : \iota$  be given. Assume the following.

$$c\_2Earithmic\_2E\_2A \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (7)$$

Let  $c\_2Earithmic\_2EDIV : \iota$  be given. Assume the following.

$$c\_2Earithmic\_2EDIV \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (8)$$

**Definition 9** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 10** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 11** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_21 2))$

**Definition 12** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))))$

**Definition 13** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A$ .if  $(\exists x \in A.p (ap P x))$  then  $(the (\lambda x.x \in A \wedge P x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 14** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40$

**Definition 15** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum$

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum.((ap (ap c\_2Earithmic\_2E\_2A (ap c\_2Earithmic\_2ENUMERAL (ap c\_2Earithmic\_2EBIT1 c\_2Earithmic\_2EZERO))) V0m) = V0m)) \quad (9)$$

Assume the following.

$$(\forall V0n \in ty\_2Enum\_2Enum.(\forall V1r \in ty\_2Enum\_2Enum.(p (ap (ap c\_2Eprim\_rec\_2E\_3C V1r) V0n)) \Rightarrow (\forall V2q \in ty\_2Enum\_2Enum.((ap (ap c\_2Earithmic\_2EDIV (ap (ap c\_2Earithmic\_2E\_2B (ap (ap c\_2Earithmic\_2E\_2A V2q) V0n)) V1r)) V0n) = V2q)))))) \quad (10)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (11)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (12)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (13)$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1x_{27} \in 2.(\forall V2y \in 2.(\forall V3y_{27} \in 2.(((p V0x) \Leftrightarrow (p V1x_{27})) \wedge ((p V1x_{27}) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_{27})))))) \Rightarrow ((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_{27}) \Rightarrow (p V3y_{27})))))) \quad (14)$$

**Theorem 1**

$$(\forall V0r \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.(p (ap (ap (ap c\_2Eprim\_rec\_2E\_3C V0r) V1n)) \Rightarrow ((ap (ap c\_2Earithmetic\_2EDIV (ap (ap c\_2Earithmetic\_2E\_2B V1n) V0r)) V1n) = (ap c\_2Earithmetic\_2ENUMERAL (ap c\_2Earithmetic\_2EBIT1 c\_2Earithmetic\_2EZERO))))))$$