

thm\_2Ebit\_2ELESS\_EQ\_EXP\_MULT  
(TMPyeK95we7nvfbVQxS6nm4kyJTVvgr3Svy)

October 26, 2020

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{1}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{2}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{3}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

**Definition 3** We define  $c\_2Earithmetic\_2EZERO$  to be  $c\_2Enum\_2E0$ .

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{4}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{5}$$

**Definition 4** We define  $c\_2Ebool\_2ET$  to be  $(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 5** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V1x \in 2.V1x))$

**Definition 6** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num\ (ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{6}$$

**Definition 7** We define  $c\_2Earithmetic\_2EBIT2$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap (ap c\_2Earithmetic\_2EBIT2 V0n) V0n)$ .

**Definition 8** We define  $c\_2Earithmetic\_2ENUMERAL$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

Let  $c\_2Earithmetic\_2E\_2A : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2A \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum} \quad (7)$$

Let  $c\_2Earithmetic\_2EEXP : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2EEXP \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum} \quad (8)$$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge p x)) \text{ of type } \iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A.\lambda P : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c\_2Emin\_2E\_40 V0P) P)))$ .

**Definition 11** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 12** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 13** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F))$ .

**Definition 14** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$ .

**Definition 15** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.V0m$ .

**Definition 16** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$ .

**Definition 17** We define  $c\_2Earithmetic\_2E\_3C\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.V0m$ .

Assume the following.

$$p (ap (ap c\_2Earithmetic\_2E\_3C\_3D V0m) (ap (ap c\_2Earithmetic\_2E\_2B V0m) V1n)))) \quad (9)$$

Assume the following.

$$(ap (ap c\_2Earithmetic\_2E\_2A V0m) V1n) = (ap (ap c\_2Earithmetic\_2E\_2A V1n) V0m)) \quad (10)$$

Assume the following.

$$(ap (ap c\_2Earithmetic\_2E\_2B V0p) V1q) = (ap (ap c\_2Earithmetic\_2E\_2A (ap (ap c\_2Earithmetic\_2EEXP V2n) V0p)) (ap (ap c\_2Earithmetic\_2EEXP V2n) V1q)))) \quad (11)$$

Assume the following.

$$\begin{aligned}
& (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. ( \\
& (p (ap (ap (ap c\_2Earithmetic\_2E\_3C\_3D V0m) V1n)) \Rightarrow (\exists V2p \in ty\_2Enum\_2Enum. \\
& (V1n = (ap (ap c\_2Earithmetic\_2E\_2B V0m) V2p))))))
\end{aligned} \tag{12}$$

Assume the following.

$$True \tag{13}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \tag{14}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\
& (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg( \\
& p V0t))))))
\end{aligned} \tag{15}$$

**Theorem 1**

$$\begin{aligned}
& (\forall V0a \in ty\_2Enum\_2Enum. (\forall V1b \in ty\_2Enum\_2Enum. ( \\
& (p (ap (ap (ap c\_2Earithmetic\_2E\_3C\_3D V0a) V1b)) \Rightarrow (\exists V2p \in ty\_2Enum\_2Enum. \\
& ((ap (ap c\_2Earithmetic\_2EEXP (ap c\_2Earithmetic\_2ENUMERAL ( \\
& ap c\_2Earithmetic\_2EBIT2 c\_2Earithmetic\_2EZERO))) V1b) = (ap \\
& (ap c\_2Earithmetic\_2E\_2A V2p) (ap (ap c\_2Earithmetic\_2EEXP (ap \\
& c\_2Earithmetic\_2ENUMERAL (ap c\_2Earithmetic\_2EBIT2 c\_2Earithmetic\_2EZERO))) \\
& V0a))))))
\end{aligned}$$