

thm_2Ebit_2ENOT__BITS2 (TMZqUMQiC7wX4UK6DkPejSHDQLrYMkNsRGP)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o\ (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1P \in 2.V1P))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (c_2Enum_2ESUC_REP\ m))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 7 We define $c_2Earithmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmetic_2EBIT1 V0n) V0n)$

Definition 8 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Definition 9 We define $c_2Earithmetic_2EBIT2$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmetic_2EBIT2 V0n) V0n)$

Let $c_2Earithmetic_2EEXP : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EEXP \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (7)$$

Let $c_2Earithmetic_2EDIV : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EDIV \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (8)$$

Definition 10 We define $c_2Ebit_2EDIV_2EXP$ to be $\lambda V0x \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.V1n$

Let $c_2Earithmetic_2E_2D : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2D \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (9)$$

Let $c_2Earithmetic_2EMOD : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EMOD \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (10)$$

Definition 11 We define $c_2Ebit_2EMOD_2EXP$ to be $\lambda V0x \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.V1n$

Definition 12 We define c_2Ebit_2EBITS to be $\lambda V0h \in ty_2Enum_2Enum.\lambda V1l \in ty_2Enum_2Enum.V1l$

Definition 13 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21) 2) (\lambda V0t \in 2.V0t)$.

Definition 14 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 15 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_21) V0t)$

Definition 16 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21) 2) (\lambda V2t \in 2.V2t) V1t2) V0t1)$

Assume the following.

$$\begin{aligned} & (\forall V0n \in ty_2Enum_2Enum.(\forall V1a \in ty_2Enum_2Enum. \\ & (\neg((ap (ap (ap c_2Ebit_2EBITS V0n) V0n) V1a) = c_2Enum_2E0)) \Leftrightarrow ((\\ & ap (ap (ap c_2Ebit_2EBITS V0n) V0n) V1a) = (ap c_2Earithmetic_2ENUMERAL \\ & (ap c_2Earithmetic_2EBIT1 c_2Earithmetic_2EZERO)))))) \end{aligned} \quad (11)$$

Assume the following.

$$True \quad (12)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p \ V0t))) \Leftrightarrow (p \ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (13)$$

Assume the following.

$$\forall A_27a.nonempty \ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (14)$$

Assume the following.

$$\forall A_27a.nonempty \ A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (15)$$

Theorem 1

$$\begin{aligned} & (\forall V0n \in ty_2Enum_2Enum.(\forall V1a \in ty_2Enum_2Enum.(\\ & (\neg((ap \ (ap \ (ap \ c_2Ebit_2EBITS \ V0n) \ V0n) \ V1a) = (ap \ c_2Earithmetic_2ENUMERAL \\ & (ap \ c_2Earithmetic_2EBIT1 \ c_2Earithmetic_2EZERO)))) \Leftrightarrow ((ap \ (\\ & ap \ (ap \ c_2Ebit_2EBITS \ V0n) \ V0n) \ V1a) = c_2Enum_2E0)))) \end{aligned}$$