

thm\_2Ebit\_2ENOT\_ZERO\_ADD1  
(TMWPeB72r6JiYzkMFSUnzj2Zekzwj6XGkY6)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.^{27a} : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a})))$

**Definition 5** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{1}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{2}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{3}$$

**Definition 6** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

**Definition 7** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 8** We define  $c\_2Ebool\_2E\_2E$  to be  $(\lambda V0t \in 2.(ap (ap\ c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E\_2F))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{4}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{5}$$

**Definition 9** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num$

**Definition 10** We define  $c\_2Earithmetic\_2EZERO$  to be  $c\_2Enum\_2E0$ .

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (6)$$

**Definition 11** We define  $c\_2Earithmetic\_2EBIT1$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap\ (ap\ c\_2Earithmetic$

**Definition 12** We define  $c\_2Earithmetic\_2ENUMERAL$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

**Definition 13** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.$ if  $(\exists x \in A.p\ (ap\ P\ x))$  then (the  $(\lambda x.x \in A \wedge$

**Definition 14** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 15** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum$

Assume the following.

$$\begin{aligned} & ((\forall V0n \in ty\_2Enum\_2Enum.((ap\ (ap\ c\_2Earithmetic\_2E\_2B \\ & \quad c\_2Enum\_2E0) V0n) = V0n)) \wedge (\forall V1m \in ty\_2Enum\_2Enum.(\forall V2n \in \\ & \quad ty\_2Enum\_2Enum.((ap\ (ap\ c\_2Earithmetic\_2E\_2B\ (ap\ c\_2Enum\_2ESUC \\ & \quad V1m)) V2n) = (ap\ c\_2Enum\_2ESUC\ (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V1m) \\ & \quad V2n)))))) \end{aligned} \quad (7)$$

Assume the following.

$$(\forall V0n \in ty\_2Enum\_2Enum.((\neg(V0n = c\_2Enum\_2E0)) \Leftrightarrow (p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C\ c\_2Enum\_2E0) V0n)))) \quad (8)$$

Assume the following.

$$\begin{aligned} & ((\forall V0m \in ty\_2Enum\_2Enum.((ap\ c\_2Enum\_2ESUC\ V0m) = (ap\ (ap \\ & \quad c\_2Earithmetic\_2E\_2B\ V0m)\ (ap\ c\_2Earithmetic\_2ENUMERAL\ (ap\ c\_2Earithmetic\_2EBIT1 \\ & \quad c\_2Earithmetic\_2EZERO)))))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & ((\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.( \\ & \quad (p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C\ V1n) V0m)) \Rightarrow (\exists V2p \in ty\_2Enum\_2Enum. \\ & \quad (V0m = (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V1n)\ (ap\ (ap\ c\_2Earithmetic\_2E\_2B \\ & \quad V2p)\ (ap\ c\_2Earithmetic\_2ENUMERAL\ (ap\ c\_2Earithmetic\_2EBIT1 \\ & \quad c\_2Earithmetic\_2EZERO)))))))))) \end{aligned} \quad (10)$$

Assume the following.

$$\forall A.\lambda 27a.nonempty\ A.\lambda 27a \Rightarrow (\forall V0x \in A.\lambda 27a.(\forall V1y \in A.\lambda 27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (11)$$

**Theorem 1**

$$(\forall V0m \in ty\_2Enum\_2Enum.((\neg(V0m = c\_2Enum\_2E0)) \Rightarrow (\exists V1p \in ty\_2Enum\_2Enum.(V0m = (ap\ c\_2Enum\_2ESUC\ V1p))))))$$