

thm_2Ebit_2ETWOEXP__DIVISION
(TMZVVhBzRY9o17pY2ZYRXLfdSkvs2D51i2o)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Earithmetic_2EMOD : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EMOD \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Earithmetic_2EDIV : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EDIV \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{3}$$

Let $c_2Earithmetic_2E_2A : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2A \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{5}$$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{6}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{7}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o(x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (8)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (9)$$

Definition 4 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num (ap$

Definition 7 We define $c_2Earithmetic_2EBIT2$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmetic_2E$

Definition 8 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Let $c_2Earithmetic_2EEXP : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EEXP \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (10)$$

Definition 9 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 10 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 11 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_21$

Definition 12 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t$

Definition 13 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then (the $(\lambda x.x \in A \wedge P x)$ of type $\iota \Rightarrow \iota$.

Definition 14 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40$

Definition 15 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum$

Assume the following.

$$(\forall V0n \in ty_2Enum_2Enum.((p (ap (ap c_2Eprim_rec_2E_3C c_2Enum_2E0) V0n)) \Rightarrow (\forall V1k \in ty_2Enum_2Enum.((V1k = (ap (ap c_2Earithmetic_2E_2B (ap (ap c_2Earithmetic_2E_2A (ap (ap c_2Earithmetic_2EDIV V1k) V0n)) V0n)) (ap (ap c_2Earithmetic_2EMOD V1k) V0n))) \wedge (p (ap (ap c_2Eprim_rec_2E_3C (ap (ap c_2Earithmetic_2EMOD V1k) V0n)) V0n)))))) \quad (11)$$

Assume the following.

$$\begin{aligned}
& (\forall V0n \in ty_2Enum_2Enum. (p (ap (ap c_2Eprim_rec_2E_3C c_2Enum_2E0) \\
& \quad (ap (ap c_2Earithmetic_2EEXP (ap c_2Earithmetic_2ENUMERAL (ap \\
& \quad \quad c_2Earithmetic_2EBIT2 c_2Earithmetic_2EZERO))) V0n)))) \tag{12}
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2. (((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\
& True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\
& \quad (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \tag{13}
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\
& (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg (p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(\\
& \quad p V0t)))))) \tag{14}
\end{aligned}$$

Theorem 1

$$\begin{aligned}
& (\forall V0n \in ty_2Enum_2Enum. (\forall V1k \in ty_2Enum_2Enum. (\\
& \quad V1k = (ap (ap c_2Earithmetic_2E_2B (ap (ap c_2Earithmetic_2E_2A \\
& \quad (ap (ap c_2Earithmetic_2EDIV V1k) (ap (ap c_2Earithmetic_2EEXP \\
& (ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT2 c_2Earithmetic_2EZERO))) \\
& \quad V0n))) (ap (ap c_2Earithmetic_2EEXP (ap c_2Earithmetic_2ENUMERAL \\
& \quad (ap c_2Earithmetic_2EBIT2 c_2Earithmetic_2EZERO))) V0n))) (\\
& \quad ap (ap c_2Earithmetic_2EMOD V1k) (ap (ap c_2Earithmetic_2EEXP \\
& (ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT2 c_2Earithmetic_2EZERO))) \\
& \quad V0n))))))
\end{aligned}$$