

thm\_2Ebool\_2EBOTH\_EXISTS\_AND\_THM  
(TM-  
LeX3RoKKyUfmNjWAgh2LJe6zVPAPN2Nyo)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2. \lambda Q \in 2. \text{inj\_o } (p \Rightarrow P \Rightarrow Q)$  of type  $\iota$ .

**Definition 2** We define `c_2Emin_2E_3D` to be  $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj\_o } (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define `c_2Ebool_2ET` to be  $(\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

**Definition 4** We define `c_2Ebool_2E_21` to be  $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^{A-27a}))))$

**Definition 5** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c\_2Ebool\_2E\_21 } 2) (\lambda V2t \in 2. V2t))))$

**Definition 6** We define `c_2Emin_2E_40` to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p (\text{ap } P x)))$  of type  $\iota \Rightarrow \iota$ .

**Definition 7** We define `c_2Ebool_2E_3F` to be  $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } V0P (\text{ap } (\text{c\_2Emin\_2E\_40 } A))))$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p \Rightarrow V0t1) \Rightarrow (p \Rightarrow V1t2)) \Rightarrow (((p \Rightarrow V1t2) \Rightarrow (p \Rightarrow V0t1)) \Rightarrow ((p \Rightarrow V0t1) \Leftrightarrow (p \Rightarrow V1t2)))))) \quad (1)$$

**Theorem 1**

$$\forall A. 27a. \text{nonempty } A. 27a \Rightarrow (\forall V0P \in 2. (\forall V1Q \in 2. ((\exists V2x \in A. 27a. ((p \Rightarrow V0P) \wedge (p \Rightarrow V1Q))) \Leftrightarrow ((\exists V3x \in A. 27a. (p \Rightarrow V0P)) \wedge (\exists V4x \in A. 27a. (p \Rightarrow V1Q))))))$$