

thm_2Ebool_2EBOTH_FORALL_IMP_THM
(TMT3Gb7qUZpyyN4P3aM69EMkyp2EeTUSCmA)

October 26, 2020

Definition 1 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p \text{ (ap } P \ x)) \text{ then (the } (\lambda x. x \in A \wedge p \text{ (ap } P \ x)) \text{ of type } \iota \Rightarrow \iota.$

Definition 2 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota.$

Definition 3 We define `c_2Ebool_2E_3F` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } V0P \text{ (ap (c_2Emin_2E_40 } A \text{ (ap } P \ x))$

Definition 4 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow q)$ of type $\iota.$

Definition 5 We define `c_2Ebool_2E_T` to be $(\text{ap (ap (c_2Emin_2E_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

Definition 6 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap (ap (c_2Emin_2E_3D } (2^{A-27a})$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p \ V0t1) \Rightarrow (p \ V1t2)) \Rightarrow (((p \ V1t2) \Rightarrow (p \ V0t1)) \Rightarrow ((p \ V0t1) \Leftrightarrow (p \ V1t2)))))) \quad (1)$$

Theorem 1

$$\forall A. 27a. \text{nonempty } A. 27a \Rightarrow (\forall V0P \in 2. (\forall V1Q \in 2. ((\forall V2x \in A. 27a. ((p \ V0P) \Rightarrow (p \ V1Q))) \Leftrightarrow ((\exists V3x \in A. 27a. (p \ V0P)) \Rightarrow (\forall V4x \in A. 27a. (p \ V1Q))))))$$