

thm_2Ebool_2ECOND__EXPAND__OR (TMZo3CZFYgAvUMaeBipguMcXhu2pgzFwoDa)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 5 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 6 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 7 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Definition 8 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$.

Definition 10 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.($

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \vee (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \vee True) \Leftrightarrow True) \wedge \\ & (((False \vee (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee False) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee \\ & (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \quad (2)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (3)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0t1 \in A.27a. (\forall V1t2 \in A.27a. (((ap (ap (ap (c.2Ebool.2ECOND A.27a) c.2Ebool.2ET) V0t1) V1t2) = V0t1) \wedge ((ap (ap (ap (c.2Ebool.2ECOND A.27a) c.2Ebool.2EF) V0t1) V1t2) = V1t2)))))) \quad (4)$$

Theorem 1

$$(\forall V0b \in 2. (\forall V1t1 \in 2. (\forall V2t2 \in 2. ((p (ap (ap (ap (c.2Ebool.2ECOND 2) V0b) V1t1) V2t2)) \Leftrightarrow (((p V0b) \wedge (p V1t1)) \vee ((\neg(p V0b)) \wedge (p V2t2))))))))$$