

thm\_2Ebool\_2ECONJ\_\_SYM  
(TMSS7oysRk9E7oqW6wbQWpVc6NHEMWkiScM)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.^{27a} : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 5** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (1)$$

**Theorem 1**

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \wedge (p V1t2)) \Leftrightarrow ((p V1t2) \wedge (p V0t1))))))$$