

thm_2Ebool_2EFUN__EQ__THM (TMPyLfcP- fYAYo55ZEK2xsmHYYoRqnNKKPQy)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (P \Rightarrow Q)$ of type ι .

Definition 3 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a})))$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (1)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow (\forall V0f \in (A_27b^{A_27a}).(\forall V1g \in (A_27b^{A_27a}).((\forall V2x \in A_27a.((ap V0f V2x) = (ap V1g V2x)) \Rightarrow (V0f = V1g)))))) \quad (2)$$

Theorem 1

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow (\forall V0f \in (A_27b^{A_27a}).(\forall V1g \in (A_27b^{A_27a}).((V0f = V1g) \Leftrightarrow (\forall V2x \in A_27a.((ap V0f V2x) = (ap V1g V2x))))))$$