

thm_2Ebool_2EF_IMP
(TMNN3XR3EEmohLg36zsUC2Yhom5DvxE4Q6c)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 4 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF$

Theorem 1 $(\forall V0t \in 2.((\neg(p V0t)) \Rightarrow ((p V0t) \Rightarrow False)))$.