

thm_2Ebool_2EIMP__CONG
(TMU7D3VZf9mbuARzS7M9VUY4ZDqU5X9i4dr)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 5 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (1)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (2)$$

Theorem 1

$$(\forall V0x \in 2.(\forall V1x._27 \in 2.(\forall V2y \in 2.(\forall V3y._27 \in 2.(((p V0x) \Leftrightarrow (p V1x._27)) \wedge ((p V1x._27) \Rightarrow ((p V2y) \Leftrightarrow (p V3y._27)))))) \Rightarrow (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x._27) \Rightarrow (p V3y._27))))))$$