

thm_2Ebool_2EMONO__COND (TMKzwazuZ9Q3JMtjZ4WGpG68depiGgS8JTj)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define `c_2Ebool_2ET` to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

Definition 4 We define `c_2Ebool_2E_21` to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 5 We define `c_2Ebool_2EF` to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2. V0t))$.

Definition 6 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2. V2t))$

Definition 7 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \mathbf{if} (\exists x \in A. p (ap P x)) \mathbf{then} (the (\lambda x. x \in A \wedge p (ap P x)))$ of type $\iota \Rightarrow \iota$.

Definition 8 We define `c_2Ebool_2ECOND` to be $\lambda A_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A_27a. (\lambda V2t2 \in A_27a. (ap (c_2Emin_2E_40 (A_27a))$

Assume the following.

$$\begin{aligned} & \forall A_27a. nonempty A_27a \Rightarrow (\forall V0t1 \in A_27a. (\forall V1t2 \in \\ & A_27a. (((ap (ap (ap (c_2Ebool_2ECOND A_27a) c_2Ebool_2ET) V0t1) \\ & V1t2) = V0t1) \wedge ((ap (ap (ap (c_2Ebool_2ECOND A_27a) c_2Ebool_2EF) \\ & V0t1) V1t2) = V1t2)))))) \end{aligned} \tag{1}$$

Theorem 1

$$\begin{aligned} & (\forall V0x \in 2. (\forall V1y \in 2. (\forall V2z \in 2. (\forall V3w \in \\ & 2. (\forall V4b \in 2. (((p V0x) \Rightarrow (p V1y)) \Rightarrow (((p V2z) \Rightarrow (p V3w)) \Rightarrow ((p \\ & (ap (ap (ap (c_2Ebool_2ECOND 2) V4b) V0x) V2z)) \Rightarrow (p (ap (ap (ap (c_2Ebool_2ECOND \\ & 2) V4b) V1y) V3w)))))))))) \end{aligned}$$