

# thm\_2Ebool\_2EMONO\_OR (TMVT- TNynoLek2moTRvDKcaR7iw2BKxfckMh)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2. \lambda Q \in 2. inj\_o (p \Rightarrow P \Rightarrow Q)$  of type  $\iota$ .

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A. \lambda x \in A. \lambda y \in A. inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 5** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.$

**Definition 6** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p \Rightarrow V0t1) \Rightarrow ((p \Rightarrow V1t2) \Rightarrow (p \Rightarrow V2t3))) \Leftrightarrow (((p \Rightarrow V0t1) \wedge (p \Rightarrow V1t2)) \Rightarrow (p \Rightarrow V2t3)))))) \quad (1)$$

**Theorem 1**

$$(\forall V0x \in 2. (\forall V1y \in 2. (\forall V2z \in 2. (\forall V3w \in 2. (((((p \Rightarrow V0x) \Rightarrow (p \Rightarrow V1y)) \wedge ((p \Rightarrow V2z) \Rightarrow (p \Rightarrow V3w))) \Rightarrow (((p \Rightarrow V0x) \vee (p \Rightarrow V2z)) \Rightarrow ((p \Rightarrow V1y) \vee (p \Rightarrow V3w))))))))))$$