

thm_2Ebool_2ERES_EXISTS_FALSE (TMdgsYfNZfef2XU9iw86UHZd5LQcutAGpGr)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2ET` to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A_{.27a} : \iota.(\lambda V0P \in (2^{A_{.27a}}).(ap (ap (c_2Emin_2E_3D (2^{A_{.27a}}))$

Definition 4 We define `c_2Ebool_2EF` to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define `c_2Ebool_2EIN` to be $\lambda A_{.27a} : \iota.(\lambda V0x \in A_{.27a}.(\lambda V1f \in (2^{A_{.27a}}).(ap V1f V0x)))$

Definition 6 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 7 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 8 We define `c_2Emin_2E_40` to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 9 We define `c_2Ebool_2E_3F` to be $\lambda A_{.27a} : \iota.(\lambda V0P \in (2^{A_{.27a}}).(ap V0P (ap (c_2Emin_2E_40 A_{.27a}) P)))$

Definition 10 We define `c_2Ebool_2ERES_EXISTS` to be $\lambda A_{.27a} : \iota.(\lambda V0p \in (2^{A_{.27a}}).(\lambda V1m \in (2^{A_{.27a}}).(\lambda V2x \in A_{.27a}.(ap V2x (ap V1m V0p))))$

Assume the following.

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0t \in 2.((\exists V1x \in A_{.27a}.(p V0t)) \Leftrightarrow (p V0t))) \quad (1)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0P \in (2^{A_{.27a}}).(\forall V1f \in \\ & (2^{A_{.27a}}).((p (ap (ap (c_2Ebool_2ERES_EXISTS A_{.27a}) V0P) V1f)) \Leftrightarrow \\ & (\exists V2x \in A_{.27a}.(p (ap (ap (c_2Ebool_2EIN A_{.27a}) V2x) V0P)) \wedge \\ & (p (ap V1f V2x)))))) \end{aligned} \quad (3)$$

Theorem 1

$$\forall A_{27a}.nonempty\ A_{27a} \Rightarrow (\forall V0P \in (2^{A_{27a}}).((p\ (ap\ (c_2Ebool_2ERES_EXISTS\ A_{27a})\ V0P)\ (\lambda V1x \in A_{27a}.c_2Ebool_2EF))) \Leftrightarrow\ False))$$