

# thm\_2Ebool\_2ERIGHT\_OR\_OVER\_AND (TM- bkvHGxEv1FNjPwp7hnptWkHzdE5MZtCUt)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2. \lambda Q \in 2. \text{inj\_o } (p \Rightarrow P \Rightarrow Q)$  of type  $\iota$ .

**Definition 2** We define `c_2Emin_2E_3D` to be  $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj\_o } (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define `c_2Ebool_2ET` to be  $(\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

**Definition 4** We define `c_2Ebool_2E_21` to be  $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^{A-27a}))))$

**Definition 5** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c\_2Ebool\_2E\_21 } 2)) (\lambda V2t \in 2. V2t))$

**Definition 6** We define `c_2Ebool_2E_5C_2F` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c\_2Ebool\_2E\_21 } 2)) (\lambda V2t \in 2. V2t))$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p \Rightarrow V0t1) \Rightarrow (p \Rightarrow V1t2)) \Rightarrow (((p \Rightarrow V1t2) \Rightarrow (p \Rightarrow V0t1)) \Rightarrow ((p \Rightarrow V0t1) \Leftrightarrow (p \Rightarrow V1t2)))))) \quad (1)$$

**Theorem 1**

$$(\forall V0A \in 2. (\forall V1B \in 2. (\forall V2C \in 2. (((p \Rightarrow V1B) \wedge (p \Rightarrow V2C)) \vee (p \Rightarrow V0A)) \Leftrightarrow (((p \Rightarrow V1B) \vee (p \Rightarrow V0A)) \wedge ((p \Rightarrow V2C) \vee (p \Rightarrow V0A))))))$$